



The Robin Mean Value Equation I: A Random Walk Approach to the Third Boundary Value Problem

Marta Lewicka¹ · Yuval Peres²

Received: 1 February 2021 / Accepted: 19 May 2022
© The Author(s), under exclusive licence to Springer Nature B.V. 2022

Abstract

We study the family of integral equations, called the Robin mean value equations (RMV), that are local averaged approximations to the Robin-Laplace boundary value problem (RL). When posed on $C^{1,1}$ -regular domains, we prove existence, uniqueness, equiboundedness and the comparison principle for solutions to (RMV). For the continuous right hand side of (RL), we show that solutions to (RMV) converge uniformly, in the limit of the vanishing radius of averaging, to the unique $W^{2,p}$ solution, which coincides with the unique viscosity solution of (RL). We further prove the lower bound on solutions to (RMV), which is consistent with the optimal lower bound for solutions to (RL). Our proofs employ martingale techniques, where (RMV) is interpreted as the dynamic programming principle along a suitable discrete stochastic process, interpolating between the reflecting and the stopped-at-exit Brownian walks.

Keywords Robin problem · Third boundary value problem · Oblique boundary conditions · Dynamic programming principle · Random walk · Finite difference approximations · Viscosity solutions

Mathematics Subject Classification (2010) 35J05 · 35J25 · 60G50

1 Introduction

The purpose of this paper is to study the family of integral equations, parametrised by $\epsilon \rightarrow 0+$:

$$u_\epsilon(x) = (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap \mathcal{D}} u_\epsilon(y) dy + \frac{\epsilon^2}{2(N+2)} f(x) \quad \text{for all } x \in \bar{\mathcal{D}}, \quad (\text{RMV})_\epsilon$$

✉ Marta Lewicka
lewicka@pitt.edu

Yuval Peres
yuval@yuvalperes.com

¹ Department of Mathematics, University of Pittsburgh, 139 University Place, Pittsburgh, PA 15260, USA

² Kent, OH, USA

posed on a bounded domain $\mathcal{D} \subset \mathbb{R}^N$, with a bounded, Borel function f , a positive constant γ , and s_ϵ appropriately given in Eq. 2.1. We call $(\text{RMV})_\epsilon$ the *Robin mean value equation* and view it as the approximation to the *Robin-Laplace problem* (known as the *third boundary value problem*):

$$-\Delta u = f \quad \text{in } \mathcal{D}, \quad \frac{\partial u}{\partial \bar{n}} + \gamma u = 0 \quad \text{on } \partial \mathcal{D}. \tag{RL}$$

Our analysis of $(\text{RMV})_\epsilon$ relies on its probabilistic interpretation as the dynamic programming principle along a discrete stochastic process $\{X_n^{\epsilon,x}\}_{n=0}^\infty$, which starts at a given $x \in \mathcal{D}$, samples uniformly on the truncated balls $B_\epsilon(X_n^{\epsilon,x}) \cap \mathcal{D}$, and stops with probability $\gamma s_\epsilon(X_n^{\epsilon,x})$ at each consecutive position $X_n^{\epsilon,x}$. The process accumulates values of f until $\tau^{\epsilon,x}$, whereas we define:

$$u^\epsilon(x) = \frac{\epsilon^2}{2(N+2)} \mathbb{E} \left[\sum_{n=0}^{\tau^{\epsilon,x}-1} (f \circ X_n^{\epsilon,x}) \right] = \frac{\epsilon^2}{2(N+2)} \mathbb{E} \left[\sum_{n=0}^\infty (f \circ X_n^{\epsilon,x}) \Lambda_n^{\epsilon,x} \right],$$

$$\text{with } \Lambda_n^{\epsilon,x} = \prod_{i=1}^n (1 - \gamma(s_\epsilon \circ X_{i-1}^{\epsilon,x})).$$

(DPP) $_\epsilon$

The second representation above reflects the value of an infinite horizon process, where the accumulation procedure rather than terminating at the stopping time $\tau^{\epsilon,x}$, never stops. The consecutive evaluations of f are instead weighted by the probabilities $\Lambda_n^{\epsilon,x}$ of not having stopped so far. The precise definitions of $\{X_n^{\epsilon,x}\}_{n=0}^\infty$ and $\tau^{\epsilon,x}$ will be given in Section 5.

In this paper we relate the three individual problems $(\text{RMV})_\epsilon$, (RL) and (DPP) $_\epsilon$, combining the analytical and probabilistic techniques in their study. Our approach suggests how to view more general, nonlinear operators (like Δ_p, Δ_∞) subject to oblique-type boundary conditions, through their local averaged approximations of the type $(\text{RMV})_\epsilon$. This approach has been previously successfully used in the Dirichlet and Neumann cases [1, 8, 14, 19, 23, 24].

We now summarize our main results. We work under the following basic hypotheses:

$$\left[\begin{array}{l} \text{The nonempty set } \mathcal{D} \subset \mathbb{R}^N \text{ is open, bounded, connected and of regularity } \\ \mathcal{C}^{1,1}. \text{ The function } f : \bar{\mathcal{D}} \rightarrow \mathbb{R} \text{ is bounded and Borel. The coefficient } \gamma > 0 \text{ is } \\ \text{a positive constant.} \end{array} \right. \tag{BH}$$

Recall that \mathcal{D} being $\mathcal{C}^{1,1}$ regular signifies that $\partial \mathcal{D}$ is locally a graph of a $\mathcal{C}^{1,1}$ function, which is equivalent to the uniform (two-sided) supporting sphere condition; see Lemma 2.2 for details.

Theorem 1.1 *Assume (BH) and let $\epsilon \ll 1$.*

- (i) *Each problem $(\text{RMV})_\epsilon$ has a unique solution $u_\epsilon = u^\epsilon$, coinciding with the value of (DPP) $_\epsilon$, that is Borel, bounded with a bound independent of ϵ , and obeys the comparison principle. For f is continuous / Hölder continuous / Lipschitz, u_ϵ inherits the same regularity.*
- (ii) *When $f \in \mathcal{C}(\bar{\mathcal{D}})$, then $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ converges uniformly on $\bar{\mathcal{D}}$ to $u \in \mathcal{C}(\bar{\mathcal{D}})$ that is the unique viscosity solution to (RL). In fact, u coincides with the unique $W^{2,p}((D))$ solution to (RL). Since the range of p covers $(1, \infty)$, it follows that $u \in \mathcal{C}^{1,\alpha}(\bar{\mathcal{D}})$ for any $\alpha \in (0, 1)$.*

In a companion paper [16] we show that $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ converges uniformly on $\bar{\mathcal{D}}$ to the unique $W^{2,p}(\mathcal{D})$ solution to (RL), for any bounded, Borel right hand side function f . To this end, in [16] we use probability techniques involving various couplings of random walks and yielding approximate Hölder regularity of u^ϵ in $(\text{DPP})_\epsilon$ (Lipschitz in the interior and $C^{0,\alpha}$ up to the boundary of \mathcal{D} , for any $\alpha \in (0, 1)$). In the present paper, it suffices to combine the simpler martingale arguments close to the boundary of \mathcal{D} , with estimates on the Taylor remainder term for the Newtonian potential, via another probabilistic interpretation of u_ϵ , parallel to that in $(\text{DPP})_\epsilon$.

By further martingale techniques we deduce the lower bound on u^ϵ in the general case of nonnegative bounded f , in function of γ and the radius of the inner supporting balls at $\partial\mathcal{D}$:

Theorem 1.2 Assume (BH) with $f \geq 0$ and let $r > 0$ satisfy:

$$\text{for every } x \in \partial\mathcal{D} \text{ exists } B_r(a) \subset \mathcal{D} \text{ such that } |x - a| = r.$$

Then the solution to $(\text{RMV})_\epsilon$ obeys the following bound, for any radius $\bar{r} < r$ provided that $\epsilon \ll 1$:

$$u^\epsilon(x_0) \geq \frac{\bar{r}}{\gamma N} \cdot \inf_{\mathcal{D}} f \quad \text{for all } x_0 \in \bar{\mathcal{D}}.$$

Clearly, uniform convergence of $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ to u implies that $u \geq \frac{r}{\gamma N} \inf_{\bar{\mathcal{D}}} f$. This bound is optimal and equivalent to Theorem 1.2. For solutions to (RL), it may be obtained directly via the maximum principle.

1.1 Prior Results in Related Contexts

To put $(\text{RMV})_\epsilon$ in a more familiar setting, rewrite:

$$u_\epsilon(x) = p_\epsilon(x) \left(\int_{B_\epsilon(x)} \mathbb{1}_{\mathcal{D}}(y) u_\epsilon(y) dy + \frac{\epsilon^2}{2(N+2)} f(x) \right) + (1 - p_\epsilon(x)) \left(\int_{B_\epsilon(x) \cap \mathcal{D}} u_\epsilon(y) dy + \frac{\epsilon^2}{2(N+2)} f(x) \right), \quad p_\epsilon(x) = \gamma s_\epsilon(x) \cdot \frac{|B_\epsilon(x)|}{|B_\epsilon(x) \setminus \mathcal{D}|}.$$

Note that the interpolation weight p_ϵ is of order ϵ , for any $\gamma > 0$. This rests in agreement with the fact that the Neumann averaging, corresponding to a higher order operator, must prevail. Indeed, we observe that the above formula interpolates between the dynamic programming principles for:

- (i) the Dirichlet problem: $-\Delta u = f$ in \mathcal{D} , $u = 0$ on $\partial\mathcal{D}$, when $p_\epsilon = 1$. This heuristically corresponds to the limiting case $\gamma = \infty$. The probabilistic interpretation for the Dirichlet problem is classical and has been extensively studied in the literature both in the present linear case of Δ and the random walk [10], as well as in the nonlinear cases of Δ_p , $p \in (1, \infty]$ and the tug-of-war games with noise [14, 19, 23, 24]. In the continuous setting, the Perron solution to the homogeneous problem: $-\Delta u = 0$ in \mathcal{D} , $u = F$ on $\partial\mathcal{D}$ has the well known representation [10]:

$$u(x) = \mathbb{E}_x[F \circ \mathcal{B}_\tau] \doteq \mathbb{E}[F \circ (x + \mathcal{B}_{\tau_x})],$$

where $\{\mathcal{B}_t\}_{t \geq 0}$ denotes the standard N -dimensional Brownian motion, and where for each $x \in \mathcal{D}$ the stopping time τ_x is defined by: $\tau_x = \min\{t \geq 0; x + \mathcal{B}_t \in \partial\mathcal{D}\}$.

- (ii) the Neumann problem: $-\Delta u = f$ in \mathcal{D} , $\frac{\partial u}{\partial n} = 0$ on $\partial\mathcal{D}$, when $p_\epsilon = 0$. This corresponds to the case $\gamma = 0$ which is not covered in our paper. Indeed, the Neumann problem is not well posed unless $\int_{\mathcal{D}} f = 0$ and most of our proofs do not work in this

limiting case. A related probabilistic approach [1] involves the reflected Brownian motion and the local boundary time, first introduced in [7] for C^3 domains. More precisely, in the continuous setting one defines the following family (nondecreasing in $t \geq 0$) of random variables, with the help of the reflected Brownian motion $\{\bar{B}_t\}_{t \geq 0}$:

$$L_t = \lim_{\delta \rightarrow 0} \frac{1}{2\delta} \int_0^t \mathbb{1}_{\{\text{dist}(\bar{B}_s, \partial\mathcal{D}) < \delta\}} \, ds.$$

Then, the solution to the homogeneous problem: $-\Delta u = 0$ in \mathcal{D} , $\frac{\partial u}{\partial \bar{n}} = F$ on $\partial\mathcal{D}$ satisfying $\int_{\partial\mathcal{D}} F = 0$, is given by the limiting expectation of the Lebesgue-Stieltjes integral in:

$$u(x) = \lim_{t \rightarrow \infty} \mathbb{E}_x \left[\int_0^t F \circ \bar{B}_s \, dL_s \right].$$

This approach was extended to $C^{2,\alpha}$ domains [4] and to Lipschitz domains [3], together with a probabilistic proof of Hölder continuity in [3, Corollary 3.8].

We finally remark that the homogeneous Robin problem for the Laplace equation: $-\Delta u = 0$ in \mathcal{D} , $\frac{\partial u}{\partial \bar{n}} + \gamma u = F$ on $\partial\mathcal{D}$, was first studied probabilistically in [22] for C^3 domains. The corresponding representation via the boundary local time of reflected Brownian motion was shown to be:

$$u(x) = \lim_{t \rightarrow \infty} \mathbb{E}_x \left[\int_0^t e^{-\gamma L_s} (F \circ \bar{B}_s) \, dL_s \right] = \mathbb{E}_x \left[\int_0^\infty e^{-\gamma L_s} (F \circ \bar{B}_s) \, dL_s \right].$$

Observe that the coefficient $e^{-\gamma L_s}$ above plays the same role as the factor $\Lambda_{[s/n]}^{\epsilon, x}$ in our discrete representation $(\text{DPP})_\epsilon$. The same problem was studied in certain fractal domains in [2]. The mixed Dirichlet-Neumann problem for Δ_∞ in the context of the game from [23], under the assumption of C^1 regular Neumann boundary and Lipschitz Dirichlet data, was addressed in [8].

1.2 Relation Between $(\text{RMV})_\epsilon$ and (RL)

To motivate the role of the coefficient $s_\epsilon(x)$ that will be precisely defined in (2.1), we average on the truncated ball $B_\epsilon(x) \cap \mathcal{D}$ the Taylor expansion:

$$u(y) = u(x) + \langle \nabla u(x), y - x \rangle + \frac{1}{2} \langle \nabla^2 u(x) : (y - x)^{\otimes 2} \rangle + o(|y - x|^2)$$

When $d = \text{dist}(x, \partial\mathcal{D}) \geq \epsilon$, this procedure leads to the familiar formula:

$$u(x) = \int_{B_\epsilon(x)} u(y) \, dy - \frac{\epsilon^2}{2(N+2)} \Delta u(x) + o(\epsilon^2),$$

whose leading order terms readily coincide with $(\text{RMV})_\epsilon$ after replacing $-\Delta u$ with f and after setting $s_\epsilon(x) = 0$. In case of $d < \epsilon$ when $x \approx \bar{x} \in \partial\mathcal{D}$, the same reasoning requires calculating the possibly nonzero average $\int_{B_\epsilon(x) \cap \mathcal{D}} y - x \, dy$. For sufficiently regular domains \mathcal{D} , one can approximate this term by the average on the ball $B_\epsilon(x)$ truncated with the tangent plane to $\partial\mathcal{D}$ at \bar{x} , rather than by the surface $\partial\mathcal{D}$. This simpler average may be then directly computed as: $-s_\epsilon(x) \bar{n}(\bar{x}) \sim -\epsilon \left(1 - \left(\frac{d}{\epsilon}\right)^2\right)^{\frac{N+1}{2}} \bar{n}(\bar{x})$. Assuming the boundary condition $u(\bar{x}) + \gamma \frac{\partial u}{\partial \bar{n}}(\bar{x}) = 0$, the first two terms of the discussed averaged Taylor expansion thus become:

$$\begin{aligned} u(x) - \langle \nabla u(x), s_\epsilon(x) \bar{n}(\bar{x}) \rangle &= u(x) - s_\epsilon(x) \frac{\partial u}{\partial \bar{n}}(\bar{x}) + O(\epsilon s_\epsilon(x)) \\ &= u(x) + \gamma s_\epsilon(x) u(x) + O(\epsilon s_\epsilon(x)). \end{aligned}$$

Since $(1 + \gamma s_\epsilon)^{-1} = (1 - \gamma s_\epsilon) + O(s_\epsilon^2)$, we conclude that:

$$u(x) = (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x)} u(y) dy - \frac{\epsilon^2}{2(N + 2)} \Delta u(x) + O(\epsilon s_\epsilon(x)) + o(\epsilon^2), \quad (1.1)$$

which coincides with $(RMV)_\epsilon$ at its leading order terms.

We remark that the Robin problem (RL), called also the third boundary value problem / impedance boundary problem / convective boundary problem, has received attention due to its applications in many contexts in science and engineering. Using classical Schauder estimates, it follows [11, Chapter 6.7] that on a bounded domain \mathcal{D} of regularity $C^{2,\alpha}$, the general strictly elliptic problem $Lu = f$ with $C^\alpha(\bar{\mathcal{D}})$ -regular coefficients and $f \in C^\alpha(\bar{\mathcal{D}})$, subject to the oblique boundary conditions: $\langle \beta(x), \nabla u(x) \rangle + \gamma(x)u(x) = \phi(x)$ posed with $\gamma, \beta, \phi \in C^{1,\alpha}(\partial\mathcal{D})$ where $\gamma \langle \beta, \vec{n} \rangle > 0$, has a unique solution $u \in C^{2,\alpha}(\bar{\mathcal{D}})$, that satisfies the usual a-priori bounds.

Much of the modern theory for nonlinear boundary value problems modeled on (RL), is contained in the recent extensive monograph [17]. It is shown in Theorem 1.26 there, that solutions to linear oblique problems in Lipschitz domains are Hölder continuous. Further, in Theorem 4.40 and Corollary 4.41 it is shown that regularity $C^{1,\alpha}$ of \mathcal{D} suffices for the solution regularity $u \in C^{2,\alpha}(\bar{\mathcal{D}})$, provided that $f \in C^\alpha(\bar{\mathcal{D}})$ and $\beta \in C^{1,\alpha}(\partial\mathcal{D})$. We observe that for (RL), the obliqueness vector $\beta = \vec{n}$ is only Lipschitz and thus one cannot, in general, expect that $u \in C^{2,\alpha}(\bar{\mathcal{D}})$. For example, when $N = 2$ then in local coordinates $\partial\mathcal{D}$ may be parametrised as the graph $\{(x_1, \phi(x_1))\}$ of some function $\phi \in C^{1,1}$. Writing $\vec{n} = (\phi'(x_1), -1)/\sqrt{1 + |\phi'(x_1)|^2}$, it is not hard to check that if $u \in C^{2,\alpha}(\bar{\mathcal{D}})$, then the boundary condition in (RL) implies that ϕ' is a solution to a quadratic equation with $C^{1,\alpha}$ coefficients, and thus it must automatically be $C^{1,\alpha}$.

In [17, Theorem 6.30] it is proved that if \mathcal{D} has regularity $C^{1,\alpha}$ and $f \in L^p(\mathcal{D})$ with $p \in (1, \frac{1}{1-\alpha})$ then $u \in W^{2,p}(\mathcal{D})$ in (RL). Consequently, when $f \in L^\infty$, we get that $u \in C^{1,\alpha}(\bar{\mathcal{D}})$ for any $\alpha \in (0, 1)$. Analysis of (RL) in non-smooth domains, including sets with a rectifiable topological boundary having finite $(N - 1)$ -dimensional Hausdorff measure, can be found in [6, 10, 21].

1.3 The Structure of this Paper

In Section 2 we prove Theorem 1.1 (i) together with some preliminary geometric lemmas regarding s_ϵ . In Section 3 we derive the expansion (1.1) for $C^2(\bar{\mathcal{D}})$ solutions u to (RL) and show the uniform convergence of $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ to u in this restricted setting. In Section 4 we recall the definition of viscosity solutions to (RL) when $f \in C(\bar{\mathcal{D}})$ and prove in Theorem 4.2, that any uniform limit of $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ must automatically be a viscosity solution. In fact, viscosity solutions to (RL) are unique under the uniform supporting spheres assumption. For completeness, we reproduce the proof of this folklore statement in Lemma 9.2 in Appendix.

In Section 5 we develop the probability setup related to $(RMV)_\epsilon$, where $u_\epsilon = u^\epsilon$. We define the stochastic process $\{X_n^\epsilon\}_{n=0}^\infty$ in $(DPP)_\epsilon$ by (5.2) and sketch two procedures achieving the uniform distribution of its positions on the truncated balls $B_\epsilon(X_n^\epsilon) \cap \mathcal{D}$: via the rejection sampling, and via the Knothe-Rosenblatt rearrangement. The first easy bound on the expectation of the stopping time τ^ϵ , of the order C/ϵ^2 , is then refined in Section 8 where we prove Theorem 8.2 that serves to deduce Theorem 1.2. As a byproduct, we obtain the lower bound on the limit u in Theorem 1.1 (ii), which is independently derived in Lemma 8.4 for $u \in C^1(\bar{\mathcal{D}})$ satisfying (RL).

The second probabilistic interpretation of u_ϵ is proposed in Section 6: instead of stopping the accumulation of values f along the process $\{X_n^\epsilon\}_{n=0}^\infty$ at τ^ϵ in $(\text{DPP})_\epsilon$, the accumulation proceeds indefinitely but each $f(X_n)$ is weighted by the product of Dirichlet factors $\Lambda_n = \prod_{i=0}^{n-1} (1 - \gamma s_\epsilon(X_i))$. The main convergence statement in Theorem 1.1 (ii) is then proved in Section 7. We first estimate the remainder term in the Robin expansion (1.1) for u that is a $W^{2,p}$ solution of (RL). The estimate at the boundary of \mathcal{D} follows from an estimate on $\sum_{n=0}^\infty \Lambda_n$ in Lemma 6.2. The estimate away from $\partial\mathcal{D}$ is given in Theorem 7.1: the said remainder equals the difference of f from its convolution with a suitable probabilistic kernel. This suffices to pass to the limit when $f \in \mathcal{C}(\bar{\mathcal{D}})$; for a bounded f that is only Borel, the same argument is refined in the companion paper [16].

1.4 Notation

For $x \in \mathbb{R}^N$, $r > 0$, by $B_r(x)$ we mean the open ball centered at x and with radius r . The integral sign $\int_A f(x) dx \doteq \frac{1}{|A|} \int_A f(x) dx$ denotes the average value of the function f , on the set $A \subset \mathbb{R}^N$ with positive Lebesgue measure $|A| > 0$. The scalar product of two matrices $X, Y \in \mathbb{R}^{N \times N}$ is $\langle X : Y \rangle \doteq \text{trace}(X^T Y)$, and the tensor product of two vectors $p, q \in \mathbb{R}^N$ is $p \otimes q = pq^T \in \mathbb{R}^{N \times N}$. Given a C^1 domain $\mathcal{D} \subset \mathbb{R}^N$, we denote by $\bar{n}(x)$ the outward unit normal vector at $x \in \partial\mathcal{D}$, and by $\pi_{\partial\mathcal{D}x}$ the projection onto $\partial\mathcal{D}$ along the normal $\bar{n}(\pi_{\partial\mathcal{D}x})$, defined for each $x \in \bar{\mathcal{D}}$ with sufficiently small distance from $\partial\mathcal{D}$. Unless specified otherwise, C stands by any universal positive constant that may depend on \mathcal{D} , γ and f , but not on ϵ , x or other parameter quantities. The Landau symbols \mathcal{O} and o likewise have the same uniformity properties. By $\epsilon \ll 1$ and $C \gg 1$ we mean a ‘‘sufficiently small’’ and a ‘‘sufficiently large’’ positive number.

2 The Robin Mean Value Equation and Some Geometrical Lemmas

In this section, we prove Theorem 1.1 (i). Recall that we work under the basic hypotheses (BH) and that we are concerned with the family of equations, parametrised by $\epsilon > 0$, given by:

$$u_\epsilon(x) = (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap \mathcal{D}} u_\epsilon(y) dy + \frac{\epsilon^2}{2(N + 2)} f(x) \quad \text{for all } x \in \bar{\mathcal{D}}. \quad (\text{RMV})_\epsilon$$

To define the coefficient $s_\epsilon(x)$ above, we introduce the following notation (see Fig. 1):

$$d_\epsilon(x) = \min \left\{ 1, \frac{1}{\epsilon} \text{dist}(x, \partial\mathcal{D}) \right\} \in [0, 1] \quad \text{for all } x \in \mathcal{D}, \epsilon > 0,$$

$$B_1^k = B_1(0) \subset \mathbb{R}^k, \quad B_{1,d}^k = B_1^k \cap \{y_k < d\} \quad \text{for all } d \in [0, 1].$$

Then we set:

$$s_\epsilon(x) = \frac{|B_1^{N-1}|}{(N + 1)|B_{1,d_\epsilon(x)}^N|} \cdot \epsilon (1 - d_\epsilon(x)^2)^{\frac{N+1}{2}} \quad \text{for all } x \in \bar{\mathcal{D}}, \epsilon > 0. \quad (2.1)$$

In order to prove existence and uniqueness of solutions to $(\text{RMV})_\epsilon$, we need some geometrical lemmas regarding the quantity s_ϵ .




Fig. 1 The referential truncated ball and the scaled distances from $\partial\mathcal{D}$

Lemma 2.1 Assume that $B_\epsilon(x) \cap \mathcal{D} = x + \epsilon B_{1,d_\epsilon(x)}^N$. Then:

$$\int_{B_\epsilon(x) \cap \mathcal{D}} y - x dy = -s_\epsilon(x) e_N.$$

Proof The result follows by an explicit calculation:

$$\begin{aligned} \int_{B_\epsilon(x) \cap \mathcal{D}} y - x dy &= \epsilon \left(\int_{B_{1,d_\epsilon(x)}^N} y_N dy \right) e_N = \epsilon \frac{|B_1^{N-1}|}{|B_{1,d_\epsilon(x)}^N|} \left(\int_{-1}^{d_\epsilon(x)} s(1-s^2)^{\frac{N-1}{2}} ds \right) e_N \\ &= -\epsilon \frac{|B_1^{N-1}|}{(N+1)|B_{1,d_\epsilon(x)}^N|} (1-d_\epsilon(x)^2)^{\frac{N+1}{2}} e_N = -s_\epsilon(x) e_N, \end{aligned}$$

where we applied change of variables and the fact that $\int_{B_{1,d}^N} y_i dy = 0$ for all $i \neq N$. \square

We now extend Lemma 2.1 to domains with curved boundaries. Recall that \mathcal{D} is said to be of class $\mathcal{C}^{1,1}$ provided that $\partial\mathcal{D}$ is locally a graph of a $\mathcal{C}^{1,1}$ function. Equivalently, such \mathcal{D} satisfies the uniform (two-sided) supporting sphere condition, stated below. This result has been first shown in [18, Section 2]; we refer to [15] for a self-contained discussion and an elementary proof.

Lemma 2.2 An open, bounded set $\mathcal{D} \subset \mathbb{R}^N$ is of class $\mathcal{C}^{1,1}$ if and only if there exists a radius $r > 0$ such that for every $x \in \partial\mathcal{D}$ there exist $a, b \in \mathbb{R}^N$ satisfying:

$$B_r(a) \subset \mathcal{D}, \quad B_r(b) \subset \mathbb{R}^N \setminus \bar{\mathcal{D}} \quad \text{and} \quad |x - a| = |x - b| = r.$$

Moreover, the global Lipschitz constant of \bar{n} can be taken as the inverse of the supporting radius:

$$|\bar{n}(x) - \bar{n}(y)| \leq \frac{1}{r} |x - y| \quad \text{for all } x, y \in \partial\mathcal{D}.$$

Lemma 2.3 Let \mathcal{D} be as in (BH). Then, for all $\epsilon \ll 1$ and all $x \in \bar{\mathcal{D}}$ we have:

(i) $\frac{|B_\epsilon(x) \setminus \mathcal{D}|}{|B_\epsilon(0)|} \leq C \frac{s_\epsilon(x)}{\epsilon}.$

(ii) We bound the volume of the following symmetric difference:

$$\frac{|(B_\epsilon(x) \cap \mathcal{D}) \Delta (x + \epsilon B_{1,d_\epsilon(x)}^N)|}{|B_\epsilon(0)|} \leq C s_\epsilon(x) \quad \text{for all } x \in \bar{\mathcal{D}}, \quad (2.2)$$

where for $\text{dist}(x, \partial\mathcal{D}) < \epsilon$ we set $R_x \in SO(N)$ to be a rotation satisfying $R_x e_N = \bar{n}(\pi_{\partial\mathcal{D}}(x))$, whereas for $\text{dist}(x, \partial\mathcal{D}) \geq \epsilon$ both sides of (2.2) are null for any $R_x \in SO(N)$.

(iii) There holds:

$$\int_{B_\epsilon(x) \cap \mathcal{D}} y - x dy = -s_\epsilon(x) \bar{n}(\pi_{\partial\mathcal{D}}x) + \mathcal{O}(\epsilon s_\epsilon(x)) \quad \text{for all } x \in \bar{\mathcal{D}}. \quad (2.3)$$

Proof 1. Let $r > 0$ be the radius of the supporting spheres as in Lemma 2.2 (see Fig. 2). It suffices to treat the case in which $\text{dist}(x, \partial\mathcal{D}) < \epsilon \ll r$ and $\bar{n}(\pi_{\partial\mathcal{D}}x) = e_N$. We first prove (ii). Since $\partial\mathcal{D}$ is contained in the region between the two supporting spheres, the quantity in (2.2) is bounded by:

$$\begin{aligned} & \frac{|B_\epsilon(x) \setminus (B_r(x + (\epsilon d_\epsilon(x) - r)e_N) \cup B_r(x + (\epsilon d_\epsilon(x) + r)e_N))|}{|B_\epsilon(0)|} \\ & \leq \frac{2 \cdot |\epsilon B_{1,d_\epsilon}^N \setminus B_r((\epsilon d_\epsilon(x) - r)e_N)|}{|B_\epsilon(0)|} \leq C |B_{1,d_\epsilon}^N \setminus B_{r/\epsilon}((d_\epsilon(x) - \frac{r}{\epsilon})e_N)|. \end{aligned}$$

Writing $d = d_\epsilon(x)$ and $\bar{r} = \frac{r}{\epsilon}$, the measure above is further bounded by the following quantity:

$$q_{d,\bar{r}} = \int_{\frac{2\bar{r}d-1-d^2}{2(\bar{r}-d)}}^d (1-s^2)^{\frac{N-1}{2}} ds. \quad (2.4)$$

Observe that for $\bar{r} \geq 2$ we have: $d - \frac{2\bar{r}d-1-d^2}{2(\bar{r}-d)} = \frac{1-d^2}{2(\bar{r}-d)} \leq \frac{1}{2(\bar{r}-1)} \leq \frac{1}{\bar{r}} = \frac{\epsilon}{r}$, so automatically:

$$q_{d,\bar{r}} \leq C\epsilon \leq Cs_\epsilon(x) \quad \text{for all } d \in \left[0, \frac{1}{2}\right].$$

On the other hand, for $d \in (\frac{1}{2}, 1]$ and $\bar{r} \geq 2$, the lower integration limit in (2.4) is nonnegative, and thus we may use the fact that the function $s \mapsto (1-s^2)^{\frac{N-1}{2}}$ is decreasing




Fig. 2 The region estimated in the proof of Lemma 2.3 (ii)

on $[0, 1]$, to obtain:

$$\begin{aligned}
 q_{d,\bar{r}} &\leq C \left(d - \frac{2\bar{r}d - 1 - d^2}{2(\bar{r} - d)} \right) \left(1 - \left(\frac{2\bar{r}d - 1 - d^2}{2(\bar{r} - d)} \right)^2 \right)^{\frac{N-1}{2}} \\
 &\leq C \frac{1 - d^2}{\bar{r}} (1 - d^2)^{\frac{N-1}{2}} = C \frac{\epsilon}{r} (1 - d^2)^{\frac{N+1}{2}} \leq C s_\epsilon(x) \quad \text{for all } d \in \left(\frac{1}{2}, 1 \right),
 \end{aligned}$$

in virtue of the simple estimate:

$$1 - \left(\frac{2\bar{r}d - 1 - d^2}{2(\bar{r} - d)} \right)^2 \leq 1 - \left(d - \frac{1 - d^2}{\bar{r}} \right)^2 \leq (1 - d^2) \left(1 + \frac{2}{\bar{r}} \right).$$

This concludes the proof of (2.2), for all $\epsilon \ll 1$ that guarantee $\bar{r} \geq 2$.

2. The bound (2.3) in (iii) is implied directly by (2.2) and Lemma 2.1. To show (i), we estimate:

$$\begin{aligned}
 \frac{|B_\epsilon(x) \setminus \mathcal{D}|}{|B_\epsilon(0)|} &\leq \frac{|B_1^N \setminus B_{1,d}^N|}{|B_1^N|} + q_{d,\bar{r}} \leq C \int_d^1 (1 - s^2)^{\frac{N-1}{2}} ds + q_{d,\bar{r}} \\
 &\leq C(1 - d^2)^{\frac{N+1}{2}} + q_{d,\bar{r}} \leq C \frac{s_\epsilon(x)}{\epsilon} + C s_\epsilon(x),
 \end{aligned}$$

which ends the proof. □

The next result provides the second order counterpart of the first order bound (2.3):

Lemma 2.4 *Let \mathcal{D} be as in (BH). Then:*

$$\int_{B_\epsilon(x) \cap \mathcal{D}} (y - x)^{\otimes 2} dy = \frac{\epsilon^2}{N + 2} Id_N + \mathcal{O}(\epsilon s_\epsilon(x)) \quad \text{for all } x \in \bar{\mathcal{D}}. \quad (2.5)$$

Proof We first assume that $B_\epsilon(x) \cap \mathcal{D} = x + \epsilon B_{1,d_\epsilon(x)}^N$. By a change of variables:

$$\begin{aligned}
 \int_{B_\epsilon(x) \cap \mathcal{D}} (y - x)^{\otimes 2} dy &= \epsilon^2 \int_{B_{1,d_\epsilon(x)}^N} y^{\otimes 2} dy = \epsilon^2 \int_{B_1^N} y^{\otimes 2} dy + \epsilon^2 \mathcal{O}(|B_1^N \setminus B_{1,d_\epsilon(x)}^N|) \\
 &= \frac{\epsilon^2}{N + 2} Id_N + \mathcal{O}(\epsilon s_\epsilon(x)),
 \end{aligned}$$

where we have used that: $\int_{B_1^N} y^{\otimes 2} dy = (\int_{B_1^N} y_1^2 dy) Id_N = \frac{1}{N+2} Id_N$, and:

$$|B_1^N \setminus B_{1,d_\epsilon(x)}^N| = |B_1^{N-1}| \int_{d_\epsilon(x)}^1 (1 - s^2)^{\frac{N-1}{2}} ds \leq C(1 - d_\epsilon(x)^2)^{\frac{N+1}{2}} = C \frac{s_\epsilon(x)}{\epsilon}.$$

In the general case, we apply the bound (2.2) which introduces an additional error term of order $\mathcal{O}(\epsilon^2 s_\epsilon(x)) \leq \mathcal{O}(\epsilon s_\epsilon(x))$. This ends the proof. □

We are now ready for the main statement of this section:

Theorem 2.5 *Assume (BH). For every $\epsilon \ll 1$, the problem (RMV) $_\epsilon$ has a unique solution $u_\epsilon = u_\epsilon^f$ that is bounded and Borel. These solutions obey comparison principle, in the sense that $f \leq g$ implies $u_\epsilon^f \leq u_\epsilon^g$ in $\bar{\mathcal{D}}$. The family $\{u_\epsilon^f\}_{\epsilon \rightarrow 0}$ is equibounded, with a bound independent of ϵ .*

Proof 1. For every bounded, Borel $u : \bar{D} \rightarrow \mathbb{R}$, define $T_\epsilon^f u : \bar{D} \rightarrow \mathbb{R}$ by:

$$T_\epsilon^f u(x) = (1 - \gamma_{s_\epsilon}(x)) \int_{B_\epsilon(x) \cap \bar{D}} u(y) \, dy + \frac{\epsilon^2}{2(N+2)} f(x) \quad \text{for all } x \in \bar{D}.$$

Clearly, the operator T_ϵ^f is monotone, in the sense that $u \leq v$ implies $T_\epsilon^f(u) \leq T_\epsilon^f(v)$ in \bar{D} . Assume first that $f \geq 0$ and recursively define the sequence of Borel functions $\{u_n\}_{n=1}^\infty$:

$$u_0 \equiv 0, \quad u_n = (T_\epsilon^f)^n u_0.$$

Since $u_1 \geq 0$, it follows that $\{u_n\}_{n=1}^\infty$ is pointwise increasing. Below, we will show that it is uniformly bounded. This property will automatically result in the pointwise convergence to some bounded, nonnegative, Borel limit function $u_\epsilon : \bar{D} \rightarrow \mathbb{R}$. Applying the monotone convergence theorem, we get that $u_{n+1} = T_\epsilon^f u_n$ converges to $T_\epsilon^f u_\epsilon$ and therefore $u_\epsilon = T_\epsilon^f u_\epsilon$ solves $(RMV)_\epsilon$. For a sign changing right hand side f , we decompose: $f = f^+ - f^-$ with $f^+, f^- \geq 0$ and observe that $u_\epsilon^f = u_\epsilon^{f^+} - u_\epsilon^{f^-}$ solves $(RMV)_\epsilon$ in this general case as well.

2. To achieve the claimed boundedness, we will now prove that there exist constants $C_1, C_2 \gg 1$ such that defining a quadratic function $v(x) = C_1 - C_2|x|^2$, there holds:

$$u \leq v \quad \text{implies} \quad T_\epsilon^f u \leq v \quad \text{in } \bar{D}, \tag{2.6}$$

for every bounded, Borel $u : \bar{D} \rightarrow \mathbb{R}$ and every $\epsilon \ll 1$. We start by noting that the monotonicity of T_ϵ^f yields, when C_1 is much larger than C_2 :

$$\begin{aligned} (T_\epsilon^f u - v)(x) &\leq (T_\epsilon^f v - v)(x) \\ &= \int_{B_\epsilon(x) \cap \bar{D}} v(y) - v(x) \, dy - \gamma_{s_\epsilon}(x) \int_{B_\epsilon(x) \cap \bar{D}} v(y) \, dy + \frac{\epsilon^2}{2(N+2)} f(x) \\ &\leq -C_2 \int_{B_\epsilon(x) \cap \bar{D}} |y|^2 - |x|^2 \, dy - \frac{C_1}{2} \gamma_{s_\epsilon}(x) + \mathcal{O}(\epsilon^2) \\ &\leq -2C_2 \left\langle \int_{B_\epsilon(x) \cap \bar{D}} y - x \, dy, x \right\rangle - \frac{C_1}{2} \gamma_{s_\epsilon}(x) - C_2 \int_{B_\epsilon(x) \cap \bar{D}} |y-x|^2 \, dy + \mathcal{O}(\epsilon^2), \end{aligned}$$

Recalling (2.3) and (2.5) implies that the above quantity is nonpositive, provided that we choose $C_1 \gg C_2 \gg 1$. This proves (2.6).

3. We thus conclude existence of solutions to $(RMV)_\epsilon$ and the stated comparison principle, which follows by construction. It remains to show uniqueness; to this end, consider the difference $u = u_{\epsilon,1} - u_{\epsilon,2}$ of some two solutions $u_{\epsilon,1}$ and $u_{\epsilon,2}$, to $(RMV)_\epsilon$. Since:

$$u(x) = (1 - \gamma_{s_\epsilon}(x)) \int_{B_\epsilon(x) \cap \bar{D}} u(y) \, dy \quad \text{for all } x \in \bar{D},$$

it follows that $u \in \mathcal{C}(\bar{D})$. Assume, by contradiction, that u attains a positive maximum at some $x_0 \in \bar{D}$. Then, there must be: $u(x_0) \leq (1 - \gamma_{s_\epsilon}(x_0))u(x_0)$, implying that: $s_\epsilon(x_0) = 0$. Consequently:

$$u(x_0) = \int_{B_\epsilon(x_0)} u(y) \, dy \leq u(x_0)$$

so $u = u(x_0)$ is constant in $B_\epsilon(x_0)$. Repeating this argument, we obtain that u attains its maximum at some $\bar{x}_0 \in \partial\bar{D} + B_\epsilon(0)$. This contradicts $s_\epsilon(\bar{x}_0) = 0$ and ends the proof of Theorem 2.5. □

Corollary 2.6 *In the context of Theorem 2.5, continuity, Hölder continuity or Lipschitz continuity of f on $\bar{\mathcal{D}}$ implies, respectively: continuity, Hölder continuity, or Lipschitz continuity of the unique solution u_ϵ^f to (RMV) $_\epsilon$.*

Proof Observe that the map s_ϵ is Lipschitz continuous on $\bar{\mathcal{D}}$ as the smooth function of the scaled distance d_ϵ , and $\bar{\mathcal{D}} \ni x \mapsto \int_{B_\epsilon(x) \cap \mathcal{D}} u(y) \, dy$ is Lipschitz as well in view of the boundedness of u . Similarly, the map $\bar{\mathcal{D}} \ni x \mapsto |B_\epsilon(x) \cap \mathcal{D}|$ is Lipschitz. Thus, the first term in the right hand side of (RMV) $_\epsilon$ is Lipschitz continuous, which proves the claim. \square

3 The Taylor Expansion and Convergence to Classical Solutions

In this section, we show the expansion (1.1) and prove convergence statement in Theorem 1.1 (ii) under higher regularity assumptions, pertaining to classical solutions of (RL).

Theorem 3.1 *Assume (BH). Let $u \in C^2(\bar{\mathcal{D}})$, $f \in C(\bar{\mathcal{D}})$ satisfy (RL). Then we have, as $\epsilon \rightarrow 0$:*

$$u(x) = (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap \mathcal{D}} u(y) \, dy + \frac{\epsilon^2}{2(N+2)} f(x) + \mathcal{O}(\epsilon s_\epsilon(x)) + o(\epsilon^2) \quad \text{for all } x \in \bar{\mathcal{D}},$$

where the Landau symbols \mathcal{O} and o are uniform in x , but may depend on u , f , \mathcal{D} and γ .

Proof Without loss of generality, we may take $u \in C^2(\mathbb{R}^N)$, so that the Taylor expansion of u results in the following expansion of the average, uniformly in $x \in \bar{\mathcal{D}}$:

$$\int_{B_\epsilon(x) \cap \mathcal{D}} u(y) \, dy = u(x) + \langle \nabla u(x), \int_{B_\epsilon(x) \cap \mathcal{D}} y - x \, dy \rangle + \frac{1}{2} \langle \nabla^2 u(x) : \int_{B_\epsilon(x) \cap \mathcal{D}} (y - x)^{\otimes 2} \, dy \rangle + o(\epsilon^2).$$

Recalling (2.3) and using boundary condition in (RL), the linear term above becomes:

$$\begin{aligned} \langle \nabla u(x), \int_{B_\epsilon(x) \cap \mathcal{D}} y - x \, dy \rangle &= -s_\epsilon(x) \langle \nabla u(x), \vec{n}(\pi_{\partial \mathcal{D}} x) \rangle + \mathcal{O}(\epsilon s_\epsilon(x)) \\ &= -s_\epsilon(x) \frac{\partial u}{\partial \vec{n}}(\pi_{\partial \mathcal{D}} x) + \mathcal{O}(\epsilon s_\epsilon(x)) \\ &= \gamma s_\epsilon(x) u(\pi_{\partial \mathcal{D}} x) + \mathcal{O}(\epsilon s_\epsilon(x)) \\ &= \gamma s_\epsilon(x) u(x) + \mathcal{O}(\epsilon s_\epsilon(x)). \end{aligned}$$

By (2.5) and (RL), the quadratic term becomes:

$$\begin{aligned} \langle \nabla^2 u(x) : \int_{B_\epsilon(x) \cap \mathcal{D}} (y - x)^{\otimes 2} \, dy \rangle &= \frac{\epsilon^2}{N+2} \langle \nabla^2 u(x) : Id_N \rangle + \mathcal{O}(\epsilon s_\epsilon(x)) \\ &= \frac{\epsilon^2}{N+2} \Delta u(x) + \mathcal{O}(\epsilon s_\epsilon(x)) = -\frac{\epsilon^2}{N+2} f(x) + \mathcal{O}(\epsilon s_\epsilon(x)). \end{aligned}$$

In conclusion, it follows that:

$$\int_{B_\epsilon(x) \cap \mathcal{D}} u(y) \, dy = (1 + \gamma s_\epsilon(x)) u(x) - \frac{\epsilon^2}{2(N+2)} f(x) + \mathcal{O}(\epsilon s_\epsilon(x)) + o(\epsilon^2). \quad (3.1)$$

Observing that:

$$\frac{1}{1 + \gamma s_\epsilon(x)} = 1 - \gamma s_\epsilon(x) + \mathcal{O}(s_\epsilon(x)^2) = 1 - \gamma s_\epsilon(x) + \mathcal{O}(\epsilon s_\epsilon(x))$$

and dividing (3.1) by $1 + \gamma s_\epsilon(x)$, yields the claim. \square

The statement below is shown using arguments similar to the proof of Theorem 2.5.

Theorem 3.2 *Under the assumptions of Theorem 3.1, the sequence of (unique) solutions $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ to (RMV) $_\epsilon$ converges to u , uniformly on \bar{D} .*

Proof 1. Define the sequence of positive numbers $\{a_\epsilon\}_{\epsilon \rightarrow 0}$, converging to 0 and such that the second error term in Taylor’s expansion in Theorem 3.1 satisfies: $o(\epsilon^2) \leq a_\epsilon \epsilon^2$, uniformly in $x \in \bar{D}$. For two parameters: a sufficiently small $\delta > 0$ and a large $C > \frac{N+2}{N\delta}$, define the quadratic functions:

$$v_\epsilon(x) = C(a_\epsilon(1 - \delta|x|^2) + \epsilon).$$

Clearly, $v_\epsilon > 0$ on \bar{D} if $\delta \ll 1$. Then we also have:

$$\begin{aligned} & v_\epsilon(x) - (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap D} v_\epsilon(y) \, dy \\ &= C a_\epsilon \left(1 - \delta|x|^2 - (1 - \gamma s_\epsilon(x)) + \delta(1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap D} |y|^2 \, dy \right) + C \epsilon s_\epsilon(x) \\ &= C a_\epsilon \left(s_\epsilon(x)(1 - \delta \int_{B_\epsilon(x) \cap D} |y|^2 \, dy) + \delta \int_{B_\epsilon(x) \cap D} |y|^2 - |x|^2 \, dy \right) + C \epsilon s_\epsilon(x). \end{aligned}$$

Writing $\int_{B_\epsilon(x) \cap D} |y|^2 - |x|^2 \, dy = \int_{B_\epsilon(x) \cap D} |y - x|^2 \, dy + 2(\int_{B_\epsilon(x) \cap D} y - x \, dy, x)$ and applying (2.3) and (2.5), the above quantity is bounded from below, again when $\delta \ll 1$, by:

$$C a_\epsilon \left(\frac{\gamma}{2} s_\epsilon(x) + \delta \left(\frac{\epsilon^2 N}{N+2} + \mathcal{O}(s_\epsilon(x)) \right) \right) + C \epsilon s_\epsilon(x) \geq C a_\epsilon \delta \frac{\epsilon^2 N}{N+2} + C \epsilon s_\epsilon(x) \geq a_\epsilon \epsilon^2 + C \epsilon s_\epsilon(x),$$

so that:

$$v_\epsilon(x) - (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap D} v_\epsilon(y) \, dy \geq a_\epsilon \epsilon^2 + C \epsilon s_\epsilon(x). \tag{3.2}$$

2. Consider the difference:

$$w_\epsilon = u - u_\epsilon - v_\epsilon.$$

By (3.2), the expansion in Lemma 3.1, and by the assumed identities (RMV) $_\epsilon$, we get:

$$w_\epsilon(x) - (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap D} w_\epsilon(y) \, dy \leq \mathcal{O}(\epsilon s_\epsilon(x)) + a_\epsilon \epsilon^2 - (a_\epsilon \epsilon^2 + C \epsilon s_\epsilon(x)) \leq 0,$$

if only $C \gg 1$. Similarly, defining: $\bar{w}_\epsilon = u - u_\epsilon + v_\epsilon$, it follows that:

$$\bar{w}_\epsilon(x) - (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap D} \bar{w}_\epsilon(y) \, dy \geq \mathcal{O}(\epsilon s_\epsilon(x)) - a_\epsilon \epsilon^2 + (a_\epsilon \epsilon^2 + C \epsilon s_\epsilon(x)) \geq 0.$$

The claimed result follows now by an application of Lemma 3.3 below, since:

$$\|u - u_\epsilon\|_{C(\bar{D})} \leq \|v_\epsilon\|_{C(\bar{D})} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0,$$

in view of $w_\epsilon \leq 0$ and $\bar{w}_\epsilon \geq 0$. □

Lemma 3.3 *If $u \in C(\bar{D})$ satisfies: $u(x) - (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap D} u(y) \, dy \leq 0$ for all $x \in \bar{D}$, then there must be: $u \leq 0$ in \bar{D} .*

Proof Let $x_0 \in \bar{D}$ be such that $u(x_0) = \max_{\bar{D}} u$ and that, by contradiction: $u(x_0) > 0$. Then:

$$u(x_0) \leq (1 - \gamma s_\epsilon(x_0)) \int_{B_\epsilon(x_0) \cap D} u(y) \, dy \leq (1 - \gamma s_\epsilon(x_0))u(x_0) \leq u(x_0),$$

implying that $s_\epsilon(x_0) = 0$ and that $u = u(x_0)$ is constant on $B_\epsilon(x_0)$. Iterating this argument, we produce another maximizer $\bar{x}_0 \in \mathcal{D}$ such that $u(\bar{x}_0) = \max_{\bar{\mathcal{D}}} u$ and $\text{dist}(\bar{x}_0, \partial\bar{\mathcal{D}}) < \epsilon$, contradicting $s_\epsilon(\bar{x}_0) = 0$. \square

4 Convergence to Viscosity Solutions

The purpose of this section is to show that any uniform limit of a sequence in $\{u_\epsilon\}_{\epsilon \rightarrow 0}$, must be a viscosity solution to the Robin problem (RL), provided that $f \in \mathcal{C}(\bar{\mathcal{D}})$. According to [9, Definition 7.4], we have the following definition:

Definition 4.1 Let $\mathcal{D} \subset \mathbb{R}^N$ be a \mathcal{C}^1 -regular domain and assume that $\gamma > 0$ and $f \in \mathcal{C}(\bar{\mathcal{D}})$. We say that $u \in \mathcal{C}(\bar{\mathcal{D}})$ is a viscosity solution to (RL), provided that:

(i) (viscosity sub-solution property)

$$\begin{aligned} \text{if } x \in \mathcal{D} \text{ and } (p, X) \in J_{\mathcal{D}}^{2,+}u(x) \text{ then: } & -\text{trace}X \leq f(x), \\ \text{if } x \in \partial\mathcal{D} \text{ and } (p, X) \in J_{\mathcal{D}}^{2,+}u(x) \text{ then: } & -\text{trace}X \leq f(x) \text{ or } \langle p, \vec{n}(x) \rangle + \gamma u(x) \leq 0. \end{aligned}$$

(ii) (viscosity super-solution property)

$$\begin{aligned} \text{if } x \in \mathcal{D} \text{ and } (p, X) \in J_{\mathcal{D}}^{2,-}u(x) \text{ then: } & -\text{trace}X \geq f(x), \\ \text{if } x \in \partial\mathcal{D} \text{ and } (p, X) \in J_{\mathcal{D}}^{2,-}u(x) \text{ then: } & -\text{trace}X \geq f(x) \text{ or } \langle p, \vec{n}(x) \rangle + \gamma u(x) \geq 0. \end{aligned}$$

The set of second order super-jets $J_{\mathcal{D}}^{2,+}u(x)$ consists of couples $(p, X) \in \mathbb{R}^N \times \mathbb{R}_{\text{sym}}^{N \times N}$, satisfying:

$$u(y) \leq u(x) + \langle p, y - x \rangle + \frac{1}{2} \langle X : (y - x)^{\otimes 2} \rangle + o(|y - x|^2) \text{ as } \bar{\mathcal{D}} \ni y \rightarrow x. \tag{4.1}$$

Analogously, the set of second order sub-jets is defined by: $J_{\mathcal{D}}^{2,-}u(x) = -J_{\mathcal{D}}^{2,+}(-u)(x)$.

Recall that the expression $\langle X : (y - x)^{\otimes 2} \rangle$ in (4.1) equals: $\langle X(y - x), y - x \rangle = \sum_{i,j=1 \dots N} X_{ij}(y_i - x_i)(y_j - x_j)$. The following is the main statement of this section:

Theorem 4.2 Assume (BH) and let $f \in \mathcal{C}(\bar{\mathcal{D}})$. If some sequence of solutions $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ to $(\text{RMV})_\epsilon$ converges uniformly on $\bar{\mathcal{D}}$, then the limit $u \in \mathcal{C}(\bar{\mathcal{D}})$ is a viscosity solution to (RL).

Proof 1. Fix $x \in \bar{\mathcal{D}}$ and $(p, X) \in J_{\mathcal{D}}^{2,+}u(x)$. For each $c > 0$ consider the quadratic test function:

$$\phi_c(y) = u(x) + \langle p, y - x \rangle + \frac{1}{2} \langle X : (y - x)^{\otimes 2} \rangle + \frac{c}{2} |y - x|^2.$$

Then: $u(x) = \phi_c(x)$, and $u < \phi_c$ in some neighbourhood of x in $\bar{\mathcal{D}}$ of the form: $(\bar{B}_{\epsilon_c}(x) \setminus \{x\}) \cap \bar{\mathcal{D}}$. Define the following sequence of positive numbers, decreasing to 0, as $j \rightarrow \infty$:

$$a_j = \min_{(\bar{B}_{\epsilon_c}(x) \setminus B_{1/j}(x)) \cap \bar{\mathcal{D}}} (\phi_c - u), \quad \text{for all } j = 1, 2, \dots$$

Let $\{\epsilon_j\}_{j \rightarrow \infty}$ be another sequence decreasing to 0, such that: $\|u_{\epsilon_j} - u\|_{C(\bar{\mathcal{D}})} < \frac{1}{2}a_j$. Further, let:

$$x_j \in \bar{B}_{\epsilon_j}(x) \cap \bar{\mathcal{D}} \quad \text{satisfy:} \quad (\phi_c - u_{\epsilon_j})(x_j) = \min_{\bar{B}_{\epsilon_j}(x) \cap \bar{\mathcal{D}}} (\phi_c - u_{\epsilon_j}).$$

We immediately obtain that $x_j \in \bar{B}_{1/j}(x) \cap \bar{\mathcal{D}}$, so that $x_j \rightarrow x$ as $j \rightarrow \infty$, because:

$$\begin{aligned} (\phi_c - u_{\epsilon_j})(y) &> (\phi_c - u)(y) - \frac{1}{2}a_j \geq \frac{1}{2}a_j \\ &> u(x) - u_{\epsilon_j}(x) = (\phi_c - u_{\epsilon_j})(x) \quad \text{for all } y \in (\bar{B}_{\epsilon_j}(x) \setminus B_{1/j}(x)) \cap \bar{\mathcal{D}}. \end{aligned}$$

We now write $(RMV)_\epsilon$ as:

$$\begin{aligned} f(x_j) &= \frac{2(N+2)}{\epsilon_j^2} \left(u_{\epsilon_j}(x_j) - (1 - \gamma s_{\epsilon_j}(x_j)) \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} u_{\epsilon_j}(y) \, dy \right) \\ &= \frac{2(N+2)}{\epsilon_j^2} \left((u_{\epsilon_j}(x_j) - \phi_c(x_j)) - (1 - \gamma s_{\epsilon_j}(x_j)) \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} u_{\epsilon_j} - \phi_c \, dy \right) \quad (4.2) \\ &\quad + \phi_c(x_j) - (1 - \gamma s_{\epsilon_j}(x_j)) \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} \phi_c(y) \, dy \\ &= \frac{2(N+2)}{\epsilon_j^2} (I_j + II_j). \end{aligned}$$

By the definition of x_j as a local minimizer of $\phi_c - u_{\epsilon_j}$, it follows that:

$$I_j \geq (u_{\epsilon_j}(x_j) - \phi_c(x_j)) \gamma s_{\epsilon_j}(x_j). \tag{4.3}$$

The bound on II_j will be achieved separately in the interior and the boundary cases.

2. If $x \in \mathcal{D}$, then $s_{\epsilon_j}(x_j) = 0$ for all $j \gg 1$, implying that $I_j \geq 0$ by (4.3). Consequently:

$$II_j = \phi_c(x_j) - \int_{B_{\epsilon_j}(x_j)} \phi_c(y) \, dy = -\frac{\epsilon_j^2}{2(N+2)} \Delta \phi_c(x) = -\frac{\epsilon_j^2}{2(N+2)} (\text{trace} X + cN).$$

Thus (4.2) becomes: $f(x_j) \geq -\text{trace} X - cN$. Passing to the limit with $j \rightarrow \infty$ and $c \rightarrow 0$, we obtain the condition requested in Definition 4.1 (i):

$$-\text{trace} X \leq f(x).$$

The same reasoning shows that $-\text{trace} X \geq f(x)$ for all $(p, X) \in J_D^{2,-} u(x)$.

3. Assume that $x \in \partial \mathcal{D}$. If for some subsequence we have: $\epsilon_j \leq \text{dist}(x_j, \partial \mathcal{D})$, then: $-\text{trace} X \leq f(x)$ by the same reasoning as in step 2 above. Therefore, it suffices to assume:

$$\text{dist}(x_j, \partial \mathcal{D}) = \epsilon_j d_j \quad \text{where } d_j = d_{\epsilon_j}(x_j) \in [0, 1) \quad \text{for all } j = 1, 2, \dots$$

Call $s_j = s_{\epsilon_j}(x_j)$ and rewrite (4.2) as follows:

$$\begin{aligned} f(x_j) &= \frac{2(N+2)}{\epsilon_j^2} \left((u_{\epsilon_j}(x_j) - \phi_c(x_j)) \gamma s_j + \phi_c(x_j) \gamma s_j \right) \quad (4.4) \\ &\quad - (1 - \gamma s_j) \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} \phi_c(y) - \phi_c(x_j) \, dy. \end{aligned}$$

We will assume the following condition:

$$(p, \vec{n}(x)) + \gamma u(x) > 0, \tag{4.5}$$

which, in particular, implies that the same quantity remains bounded away from 0, if $\vec{n}(x)$ is replaced by $\vec{n}(\pi_{\partial\mathcal{D}}x_j)$ and $u(x)$ by $\phi_c(x_j)$. Consequently, for some $a > 0$ there holds:

$$\phi_c(x_j) \geq 2a - \frac{1}{\gamma} \langle p, \vec{n}(\pi_{\partial\mathcal{D}}x_j) \rangle \quad \text{for all } j = 1, 2, \dots$$

and (4.4) becomes, in view of $\nabla\phi_c(x_j) = p + (X + cId_N)(x_j - x)$ and $\nabla^2\phi_c(x_j) = X + cId_N$:

$$\begin{aligned} f(x_j) \geq & \frac{2(N+2)}{\epsilon_j^2} \left(a\gamma s_j - s_j \langle p, \vec{n}(\pi_{\partial\mathcal{D}}x_j) \rangle - \langle p, \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} y - x_j \, dy \rangle \right. \\ & - \langle (X + cId_N)(x_j - x), \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} y - x_j \, dy \rangle \\ & \left. - \frac{1}{2} \langle X + cId_N : \int_{B_{\epsilon_j}(x_j) \cap \mathcal{D}} (y - x_j)^{\otimes 2} \, dy \rangle \right). \end{aligned}$$

We now use (2.3) and (2.5) to get, for all $j \gg 1$:

$$\begin{aligned} f(x_j) \geq & \frac{2(N+2)}{\epsilon_j^2} \left(a\gamma s_j - \frac{\epsilon_j^2}{2(N+2)} \langle X + cId_N : Id_N \rangle + \mathcal{O}(|x_j - x|s_j) + \mathcal{O}(\epsilon_j s_j) \right) \\ \geq & -\text{trace}X - cN. \end{aligned}$$

Passing to the limit with $j \rightarrow \infty$ and with $c \rightarrow 0$, we obtain: $-\text{trace}X \leq f(x)$, which is precisely the condition requested in Definition 4.1 (i), now pertaining to the case $x \in \partial\mathcal{D}$, in presence of (4.5). A similar reasoning yields that u is also a viscosity super-solution. \square

In fact, viscosity solutions to (RL) are unique under the uniform outer supporting sphere assumption. This statement follows from analysis in [9, Theorem 7.5] but since the linear operator $-\Delta$ does not satisfy the u -coercivity assumption (7.14) in there, we will sketch the related proof of comparison principle in Lemma 9.2 in the Appendix A. In Section 7 we will show that the entire sequence $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ converges to the unique $W^{2,p}$ solution to (RL). Thus, we independently obtain that viscosity solutions exist, are unique and coincide with the weak solutions. We remark that another proof of existence of viscosity solutions can be obtained by Perron’s method in view of the comparison principle [9, 12]. Finally, we anticipate that in [16] we show the asymptotic Hölder equicontinuity of $\{u_\epsilon\}_{\epsilon \rightarrow 0}$ generated by any Borel $f \in L^\infty(\bar{\mathcal{D}})$. Combined with a refinement of the present arguments, this yields the assertion of Theorem 1.1 (ii) and thus another independent proof of existence and uniqueness of solutions to (RL), in the more general case.

5 The First Probabilistic Interpretation of u_ϵ

We now develop the basic probability setting related to the eq. (RMV) $_\epsilon$. As anticipated in the introduction, we will show that the unique bounded, Borel solution to (RMV) $_\epsilon$ is given by the first probabilistic formula in (DPP) $_\epsilon$. The second formula will be discussed in Section 6.

1. Consider the probability space $(B_1^N, \mathcal{B}, \frac{1}{|B_1^N|} \mathcal{L}_N)$ equipped with the standard Borel σ -algebra and the normalised Lebesgue measure, and define $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ as the countable product of B_1^N augmented by the unit interval (likewise equipped with Borel σ -algebra

and Lebesgue measure):

$$\Omega_1 = (B_1^N)^{\mathbb{N}} \times (0, 1) = \{(w, b); w = \{w^j\}_{j=1}^\infty, w^j \in B_1^N \text{ for all } j \in \mathbb{N} \text{ and } b \in (0, 1)\}.$$

Further, the countable product of $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ is denoted by $(\Omega, \mathcal{F}, \mathbb{P})$, where:

$$\Omega = (\Omega_1)^{\mathbb{N}} = \{\omega = \{(w_i, b_i)\}_{i=1}^\infty; w_i = \{w_i^j\}_{j=1}^\infty, w_i^j \in B_1^N, b_i \in (0, 1) \text{ for all } i, j \in \mathbb{N}\}.$$

For each $n \in \mathbb{N}$, the probability space $(\Omega_n, \mathcal{F}_n, \mathbb{P}_n)$ is the product of n copies of $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$ and the σ -algebra \mathcal{F}_n is identified with the sub- σ -algebra of \mathcal{F} , consisting of sets $A \times \prod_{i=n+1}^\infty \Omega_1$ for all $A \in \mathcal{F}_n$. Then $\{\mathcal{F}_n\}_{n=0}^\infty$ where $\mathcal{F}_0 = \{\emptyset, \Omega\}$, is a filtration of \mathcal{F} .

- Given $\epsilon \ll 1$, define the sequence of measurable functions $\{k_i^\epsilon : \Omega \times \bar{\mathcal{D}} \rightarrow \mathbb{N} \cup \{+\infty\}\}_{i=1}^\infty$ by:

$$k_i^\epsilon(\omega, x) = \min \{k \geq 1; x + \epsilon w_i^k \in B_\epsilon(x) \cap \mathcal{D}\} \quad \text{for all } \omega \in \Omega, x \in \bar{\mathcal{D}}.$$

Since each k_i^ϵ is \mathbb{P} -a.s. finite, we further construct the sequence of vector-valued random variables $\{w_i^{\epsilon, x} : \Omega \rightarrow B_1^N\}_{i=1}^\infty$, corresponding to $\epsilon \ll 1$ and $x \in \bar{\mathcal{D}}$, by:

$$w_i^{\epsilon, x}(\omega) = w_i^{k_i^\epsilon(\omega, x)} \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \Omega.$$

The procedure of generating $w_i^{k_i^\epsilon}$ is well known under the name of rejection sampling and it has the following measure preservation property: for every ϵ, x as above, for every Borel set $F \subset B_\epsilon(x) \cap \mathcal{D}$:

$$\begin{aligned} \mathbb{P}(x + \epsilon w_i^{\epsilon, x} \in F) &= \sum_{k=1}^\infty \mathbb{P}_1(\{x + \epsilon w_i^k \in F\} \cap \{k = k_i^\epsilon\}) \\ &= \frac{|F|}{|B_\epsilon(x)|} \cdot \sum_{k=1}^\infty \left(1 - \frac{|B_\epsilon(x) \cap \mathcal{D}|}{|B_\epsilon(x)|}\right)^{k-1} = \frac{|F|}{|B_\epsilon(x) \cap \mathcal{D}|}. \end{aligned} \tag{5.1}$$

For $\epsilon \ll 1, x_0 \in \bar{\mathcal{D}}$, we recursively define the sequence of random variables $\{X_n^{\epsilon, x_0} : \Omega \rightarrow \bar{\mathcal{D}}\}_{n=0}^\infty$ (see Fig. 3):

$$X_0^{\epsilon, x_0} \equiv x_0, \quad X_n^{\epsilon, x_0}(w_1, \dots, w_n) = X_{n-1}^{\epsilon, x_0}(w_1, \dots, w_{n-1}) + \epsilon w_n^{\epsilon, X_{n-1}^{\epsilon, x_0}(w_1, \dots, w_{n-1})}. \tag{5.2}$$

Clearly, the function X_n^{ϵ, x_0} is \mathcal{F}_n -measurable and takes values in \mathcal{D} , for $n \geq 1$. Given $\gamma > 0$, define further the \mathcal{F} -measurable $\tau^{\epsilon, x_0} : \Omega \rightarrow \mathbb{N} \cup \{+\infty\}$ by:

$$\tau^{\epsilon, x_0}(\omega) = \min \{n \geq 1; b_n < \gamma s_\epsilon(X_{n-1}^{\epsilon, x_0})\}.$$

We observe that each $X_n^{\epsilon, x_0}(\omega)$ and τ^{ϵ, x_0} are jointly measurable in ω and x_0 , by the same property of k_i^ϵ . When no ambiguity arises, we will simply write $X_n^{x_0}$ and τ^{x_0} , to simplify the notation.

- The following elementary argument proves that τ^{x_0} is finite \mathbb{P} -a.e., making it a stopping time. It is easy to notice that there exists $\bar{n} \gg 1$ and $\delta > 0$ such that for every $x_0 \in \bar{\mathcal{D}}$ there is some $n \leq \bar{n}$, satisfying:

$$\mathbb{P}_{n-1}\left((w_1, \dots, w_{n-1}); X_{n-1}^{x_0}(w_1, \dots, w_{n-1}) \in \partial\mathcal{D} + B_{\epsilon/2}(0)\right) \geq \delta.$$

Indeed, given $x_0 \in \bar{\mathcal{D}}$, one may choose $\bar{x} \in \partial\mathcal{D}$ such that $\text{dist}(x_0, \partial\mathcal{D}) = |x_0 - \bar{x}|$ and keep advancing the position $X_i^{x_0}$ in the direction of \bar{x} , by random increments w^1

within a small sector of positive measure, located in $\frac{1}{2}B_1^N$ close to its boundary, until $X_i^{x_0} \in \partial\mathcal{D} + B_{\epsilon/2}(0)$. Further, since $1 \geq \gamma s_\epsilon(x) \geq C\epsilon$ for all $x \in (\partial\mathcal{D} + B_{\epsilon/2}(0)) \cap \mathcal{D}$ and $\epsilon \ll 1$, we get: $\mathbb{P}(\tau^{x_0} \leq \bar{n}) \geq (C\epsilon)^{\bar{n}}\delta = \bar{\delta} > 0$. By induction, it follows that: $\mathbb{P}(\tau^{x_0} > k\bar{n}) \leq (1 - \bar{\delta})^k$ for all $k \geq 0$, and so $\mathbb{P}(\tau^{x_0} = \infty) = 0$, completing the argument. It also follows that:

$$\mathbb{E}[\tau^{x_0}] = \sum_{i=0}^{\infty} \mathbb{P}(\tau^{x_0} > i) \leq \bar{n} \cdot \sum_{k=0}^{\infty} \mathbb{P}(\tau^{x_0} > k\bar{n}) \leq \frac{\bar{n}}{\bar{\delta}} = C_\epsilon. \tag{5.3}$$

In fact, the above constant C_ϵ is of the order $\frac{C}{\epsilon^2}$, as noted in Corollary 5.2.

4. For each $\epsilon \ll 1, x_0 \in \bar{\mathcal{D}}$ we define the random variable $F^{\epsilon, x_0}(\omega) = \sum_{i=0}^{\tau^{x_0}-1} (f \circ X_i^{x_0})(\omega)$ that is \mathbb{P} -integrable in view of Eq. 5.3, and set:

$$u^\epsilon(x_0) = \frac{\epsilon^2}{2(N+2)} \mathbb{E}[F^{\epsilon, x_0}] = \int_\Omega \sum_{i=0}^{\tau^{x_0}-1} \frac{\epsilon^2}{2(N+2)} (f \circ X_i^{x_0})(\omega) \, d\mathbb{P}(\omega). \tag{5.4}$$

Since the function $\Omega \times \bar{\mathcal{D}} \ni (\omega, x_0) \mapsto F^{x_0}(\omega)$ is measurable, in view of the same joint measurability of $X_n^{x_0}$ and τ^{x_0} , it follows that u^ϵ is a Borel function of x_0 . The following is the main observation:

Lemma 5.1 Assume (BH). For each $\epsilon \ll 1$, the function u^ϵ in Eq. 5.4 coincides with the unique bounded, Borel solution to (RMV) $_\epsilon$:

$$u^\epsilon = u_\epsilon.$$

Proof Boundedness of u^ϵ results from Eq. 5.3, because: $|u^\epsilon(x_0)| \leq \|f\|_{L^\infty(\mathcal{D})}\epsilon^2 C_\epsilon$ for every $x_0 \in \bar{\mathcal{D}}$. To check that u^ϵ satisfies (RMV) $_\epsilon$, observe that:

$$F^{x_0}((w_1, b_1), (w_2, b_2), \dots) = f(x_0) + \mathbb{1}_{b_1 \geq \gamma s_\epsilon(x_0)} \cdot \sum_{i=0}^{\tau^{X_1^{x_0}(w_1)}-1} f(X_i^{X_1^{x_0}(w_1)}((w_2, b_2), (w_3, b_3), \dots)).$$




Fig. 3 Positions of the process defined in Eq. 5.2

Consequently:

$$\begin{aligned}
 u^\epsilon(x_0) &= \frac{\epsilon^2}{2(N+2)} \mathbb{E}[F^{x_0}] \\
 &= \frac{\epsilon^2}{2(N+2)} \int_{\Omega_1} \int_{\prod_{i=2}^\infty \Omega_1} F^{x_0}((w_1, b_1), (w_2, b_2), \dots) d\left(\prod_{i=2}^\infty \mathbb{P}_1(w_i, b_i)\right) d\mathbb{P}_1(w_1, b_1) \\
 &= \frac{\epsilon^2}{2(N+2)} f(x_0) + \int_{\Omega_1} (1 - \gamma s_\epsilon(x_0)) \int_{\prod_{i=2}^\infty \Omega_1} \sum_{i=0}^{\tau^{x_1}-1} \frac{\epsilon^2}{2(N+2)} f(X_i^{x_1}) \\
 &\quad d\left(\prod_{i=2}^\infty \mathbb{P}_1\right) d\mathbb{P}_1(w_1, b_1) \\
 &= \frac{\epsilon^2}{2(N+2)} f(x_0) + (1 - \gamma s_\epsilon(x_0)) \int_{\Omega_1} u^\epsilon(X_1^{x_0}(w_1)) d\mathbb{P}_1(w_1).
 \end{aligned}$$

Changing the variable in the last integral and recalling the measure-preserving property (5.1) in:

$$\begin{aligned}
 u^\epsilon(x_0) &= \frac{\epsilon^2}{2(N+2)} f(x_0) + (1 - \gamma s_\epsilon(x_0)) \int_{\Omega_1} u^\epsilon(x_0 + \epsilon w_1^{k_\epsilon^f(w_1, x_0)}) d\mathbb{P}_1(w_1) \\
 &= \frac{\epsilon^2}{2(N+2)} f(x_0) + (1 - \gamma s_\epsilon(x_0)) \int_{B_\epsilon(x_0) \cap \mathcal{D}} u^\epsilon(y) dy,
 \end{aligned}$$

yields $u^\epsilon = u_\epsilon$ by the uniqueness of solutions to (RMV) $_\epsilon$. □

We finally deduce the following refinement of Eq. 5.3. In Section 8 we then prove a precise lower bound on $\mathbb{E}[\tau^{\epsilon, x_0}]$ in terms of $\frac{1}{\epsilon^2}$ and the parameters γ and r in (BH).

Corollary 5.2 *There exists a constant C, depending on \mathcal{D} and γ but not on ϵ , such that:*

$$\mathbb{E}[\tau^{\epsilon, x_0}] \leq \frac{C}{\epsilon^2} \quad \text{for all } x_0 \in \bar{\mathcal{D}} \quad \text{and all } \epsilon \ll 1.$$

Proof We have:

$$\mathbb{E}[\tau^{\epsilon, x_0}] - 1 = \frac{2(N+2)}{\epsilon^2} u_\epsilon^f(x_0) \leq \frac{C}{\epsilon^2},$$

where we used Lemma 5.1 and Theorem 2.5 applied to the constant function $f = 1$. □

Remark 5.3 Another way of defining the process in Eq. 5.2 is based on the quantile regression procedure, as we now describe. We first redefine the probability spaces, by setting $\Omega_1 = B_1^N \times (0, 1)$ and taking \mathcal{F}_1 to be the Borel σ -algebra and \mathbb{P}_1 the normalised Lebesgue measure on Ω_1 . The probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is given as the countable product of $(\Omega_1, \mathcal{F}_1, \mathbb{P}_1)$, where:

$$\Omega = (\Omega_1)^\mathbb{N} = \{\omega = \{(w_i, b_i)\}_{i=1}^\infty; w_i \in B_1^N, b_i \in (0, 1) \text{ for all } i \in \mathbb{N}\}.$$

For each $n \in \mathbb{N}$, we have the corresponding $(\Omega_n, \mathcal{F}_n, \mathbb{P}_n)$ as before. For $x_0 \in \bar{\mathcal{D}}, \epsilon \ll 1$, we now inductively define the sequence of random variables: $\{X_n^{\epsilon, x_0} : \Omega \rightarrow \bar{\mathcal{D}}\}_{n=0}^\infty$:

$$X_0^{\epsilon, x_0} \equiv x_0, \quad X_n^{\epsilon, x_0}(w_1, \dots, w_n) = X_{n-1}^{\epsilon, x_0}(w_1, \dots, w_{n-1}) + T_\epsilon(X_{n-1}^{\epsilon, x_0}, \epsilon w_n), \quad (5.5)$$

where the transformation T_ϵ is given by the classical Knothe-Rosenblatt rearrangement [13, 25], whose construction we sketch below. Each X_n^{ϵ, x_0} in Eq. 5.5 is \mathcal{F}_n -measurable and it takes values in $\bar{\mathcal{D}}$. Given $\gamma > 0$, let:

$$\tau^{\epsilon, x_0}(\omega) = \min \{n \geq 1; b_n < \gamma s_\epsilon(X_{n-1}^{\epsilon, x_0})\} \quad \text{for all } \omega \in \Omega,$$

and observe, as in Eq. 5.3 that it is a stopping time. With the same definition of u^ϵ as in Eq. 5.4, the statement in Lemma 5.1 is valid as before.

Lemma 5.4 *Let $\mathcal{D} \subset \mathbb{R}^N$ be an open, bounded, \mathcal{C}^1 -regular domain. For all $\epsilon \ll 1$ there exists a continuous transformation $T_\epsilon : \bar{\mathcal{D}} \times B_\epsilon(0) \rightarrow \mathbb{R}^N$ such that for every $x \in \bar{\mathcal{D}}$, the map $T_\epsilon(x, \cdot) : B_\epsilon(0) \rightarrow B_\epsilon(x) \cap \mathcal{D}$ is a homeomorphism that is normalised-volume preserving, i.e.:*

$$\frac{|T_\epsilon(x, F)|}{|B_\epsilon(x) \cap \mathcal{D}|} = \frac{|F|}{|B_\epsilon(0)|} \quad \text{for every Borel set } F \subset B_\epsilon(0).$$

Proof Fix $x_0 \in \bar{\mathcal{D}}$. For $\text{dist}(x_0, \partial\mathcal{D}) \geq \epsilon$, we set $T_\epsilon(x_0, \cdot) = id_N$. When $\text{dist}(x_0, \partial\mathcal{D}) < \epsilon \ll 1$ then $\partial\mathcal{D} \cap B_\epsilon(x_0)$ is a \mathcal{C}^1 graph over the hyperplane perpendicular to \vec{n} , and by rescaling and rotating, we reduce to the case:

$$x_0 = 0, \quad \epsilon = 1, \quad \vec{n} = e_N, \quad B_\epsilon(0) \cap \mathcal{D} = B_1(0) \cap \{x_N < \alpha(x_1, \dots, x_{N-1})\}.$$

The Knothe-Rosenblatt rearrangement map is best described by denoting:

$$\bar{\alpha} = \frac{\mathbb{1}_{B_1^N \cap \{x_N < \alpha(x_1, \dots, x_{N-1})\}}}{|B_1^N \cap \{x_N < \alpha(x_1, \dots, x_{N-1})\}|}, \quad \bar{\beta} = \frac{\mathbb{1}_{B_1^N}}{|B_1^N|}$$

and seeking the measure-preserving diffeomorphism $R = T_1(x_0, \cdot)$:

$$R : (B_1^N, \bar{\beta} \, dx) \rightarrow (B_1^N \cap \{x_N < \alpha\}, \bar{\alpha} \, dx),$$

of the form: $R(x_1, \dots, x_N) = (R^1(x_1), R^2(x_1, x_2), \dots, R^N(x_1, \dots, x_N))$ (see Fig. 4). The continuous and strictly increasing components $R^i(x_1, \dots, x_{i-1}, \cdot)$ are recursively defined by implicit formulas:

$$\int_{-\infty}^{R^1(x_1)} \int_{\mathbb{R}^{N-1}} \bar{\alpha}(y_1, y_2, \dots, y_N) \, d(y_2, \dots, y_N) \, dy_1 = \int_{-\infty}^{x_1} \int_{\mathbb{R}^{N-1}} \bar{\beta}(y_1, y_2, \dots, y_N) \, d(y_2, \dots, y_N) \, dy_1$$

and, for all $x_1 \in \mathbb{R}$:

$$\begin{aligned} & \int_{-\infty}^{R^2(x_1, x_2)} \int_{\mathbb{R}^{N-2}} \bar{\alpha}(R^1(x_1), y_2, \dots, y_N) \, d(y_3, \dots, y_N) \, dy_2 \\ &= \frac{\int_{\mathbb{R}^{N-1}} \bar{\alpha}(R^1(x_1), y_2, \dots, y_N) \, d(y_2, \dots, y_N)}{\int_{\mathbb{R}^{N-1}} \bar{\beta}(x_1, y_2, \dots, y_N) \, d(y_2, \dots, y_N)} \cdot \int_{-\infty}^{x_2} \int_{\mathbb{R}^{N-2}} \bar{\beta}(x_1, y_2, \dots, y_N) \, d(y_3, \dots, y_N) \, dy_2, \end{aligned}$$

eventually followed by the identity:

$$\begin{aligned} & \int_{-\infty}^{R^N(x_1, \dots, x_N)} \int_{\mathbb{R}^{N-2}} \bar{\alpha}(R^1(x_1), R^2(x_1, x_2), \dots, R^{N-1}(x_1, \dots, x_{N-1}, y_N)) \, dy_N \\ &= \frac{\int_{\mathbb{R}} \bar{\alpha}(R^1(x_1), R^2(x_1, x_2), \dots, R^{N-1}(x_1, \dots, x_{N-1}, y_N)) \, dy_N}{\int_{\mathbb{R}} \bar{\beta}(x_1, \dots, x_{N-1}, y_N) \, dy_N} \cdot \int_{-\infty}^{x_N} \bar{\beta}(x_1, \dots, x_{N-1}, y_N) \, dy_N, \end{aligned}$$

whose validity is requested for all $(x_1, \dots, x_{N-1}) \in \mathbb{R}^{N-1}$. The fact that R is measure preserving is then classical and may be checked directly. Continuity of R and of its inverse R^{-1} is implied by the property that sections of both B_1^N and $B_1^N \cap \{x_N < \alpha\}$, obtained by

fixing (x_1, \dots, x_i) , are connected sets, for any $i \leq N - 1$. Finally, $R \equiv id_N$ when $\alpha = 1$ and the construction is continuous with respect to \vec{n} , justifying continuity of T_ϵ in x_0 . \square

6 The Second Probabilistic Interpretation of u_ϵ

The second probabilistic interpretation of solutions to of $(RMV)_\epsilon$ as anticipated in $(DPP)_\epsilon$, will be given below. For each $\epsilon \ll 1$, $x_0 \in \bar{\mathcal{D}}$, we define the random variables $\{\Lambda_n^{\epsilon, x_0} : \Omega \rightarrow \mathbb{R}\}_{n=0}^\infty$ along the process $\{X_n^{\epsilon, x_0}\}_{n=0}^\infty$ in Eq. 5.2, by taking the following products:

$$\Lambda_n^{\epsilon, x_0}(\omega) = \prod_{j=1}^n (1 - \gamma(s_\epsilon \circ X_{j-1}^{\epsilon, x_0})(\omega)). \tag{6.1}$$

For $n = 0$ we adopt the convention that $\Lambda_0 = 1$. When no ambiguity arises we write $\Lambda_n^{x_0}$ or Λ_n , and note that each $\Lambda_n^{x_0}(\omega)$ is jointly measurable in ω and x_0 . We now set:

$$\begin{aligned} \bar{u}^\epsilon(x_0) &= \frac{\epsilon^2}{2(N+2)} \mathbb{E} \left[\sum_{i=0}^\infty (f \circ X_i^{x_0}) \Lambda_i^{x_0} \right] \\ &= \int_\Omega \sum_{i=0}^\infty \frac{\epsilon^2}{2(N+2)} (f \circ X_i^{x_0})(\omega) \cdot \prod_{j=1}^i (1 - \gamma(s_\epsilon \circ X_{j-1}^{\epsilon, x_0})(\omega)) \, d\mathbb{P}(\omega). \end{aligned} \tag{6.2}$$

Lemma 6.1 Assume (BH). For each $\epsilon \ll 1$, the function \bar{u}^ϵ is well defined a.e. in $\bar{\mathcal{D}}$. After a possible adjustment on a negligible set, it coincides with unique bounded Borel solution to $(RMV)_\epsilon$:

$$\bar{u}^\epsilon = u_\epsilon.$$

Proof 1. Denote by v_ϵ the solution to $(RMV)_\epsilon$ with $f \equiv 1$. We first argue that the following sequence of random variables $\{M_n\}_{n=0}^\infty$ is a martingale with respect to the filtration $\{\mathcal{F}_n\}_{n=0}^\infty$:

$$M_n = (v_\epsilon \circ X_n) \Lambda_n + \frac{\epsilon^2}{2(N+2)} \sum_{i=0}^{n-1} \Lambda_i,$$




Fig. 4 The Knothe-Rosenblatt rearrangement in Lemma 5.4

where we adopt the convention that $M_0 = v_\epsilon(x_0)$. Indeed, $(RMV)_\epsilon$ yields:

$$\begin{aligned} \mathbb{E}(M_{n+1} - M_n \mid \mathcal{F}_n) &= \left(\int_{B_\epsilon(X_n) \cap \mathcal{D}} v_\epsilon(y) \, dy \right) \cdot \Lambda_{n+1} - v_\epsilon(X_n) \cdot \Lambda_n + \frac{\epsilon^2}{2(N+2)} \Lambda_n \\ &= \left((1 - \gamma s_\epsilon(X_n)) \int_{B_\epsilon(X_n) \cap \mathcal{D}} v_\epsilon(y) \, dy - v_\epsilon(X_n) + \frac{\epsilon^2}{2(N+2)} \right) \Lambda_n \\ &= 0 \quad \mathbb{P} - \text{a.s. in } \Omega. \end{aligned}$$

Using equiboundedness of $\{v_\epsilon\}_{\epsilon \rightarrow 0}$ in Theorem 2.5 we get:

$$\frac{\epsilon^2}{2(N+2)} \mathbb{E} \left[\sum_{i=0}^n \Lambda_i \right] \leq \mathbb{E}[M_{n+1}] = \mathbb{E}[M_0] = v_\epsilon(x_0) \leq C. \tag{6.3}$$

The above uniform (independent of x_0, ϵ and n) bound suffices to conclude that the function $\Omega \times \bar{\mathcal{D}} \ni (\omega, x_0) \mapsto \frac{\epsilon^2}{2(N+2)} \sum_{i=0}^\infty (f \circ X_i^{x_0})(\omega) \Lambda_i^{x_0}(\omega)$ is jointly integrable. Consequently, \bar{u}^ϵ is well defined for a.e. $x_0 \in \bar{\mathcal{D}}$, Borel regular and bounded, and moreover:

$$\mathbb{E} \left[\sum_{i=0}^\infty (f \circ X_i^{x_0}) \Lambda_i^{x_0} \right] = \sum_{i=0}^\infty \mathbb{E} \left[(f \circ X_i^{x_0}) \Lambda_i^{x_0} \right] \tag{6.4}$$

2. Similarly as in Lemma 5.1, Fubini’s theorem and change of variable in view of Eq. 5.1 yield:

$$\begin{aligned} \bar{u}^\epsilon(x_0) &= \frac{\epsilon^2}{2(N+2)} \left(f(x_0) + (1 - \gamma s_\epsilon(x_0)) \int_{\Omega_1} \mathbb{E} \left[\sum_{i=0}^\infty (f \circ X_i^{X_1}) \cdot \prod_{j=1}^i (1 - \gamma s_\epsilon(X_{j-1}^{X_1})) \right] \, d\mathbb{P}_1 \right) \\ &= \frac{\epsilon^2}{2(N+2)} f(x_0) + (1 - \gamma s_\epsilon(x_0)) \int_{\Omega_1} \bar{u}^\epsilon(X_1^{x_0}(w_1)) \, d\mathbb{P}_1(w_1) \\ &= \frac{\epsilon^2}{2(N+2)} f(x_0) + (1 - \gamma s_\epsilon(x_0)) \int_{B_\epsilon(x_0) \cap \mathcal{D}} \bar{u}^\epsilon(y) \, dy, \end{aligned}$$

for a.e. $x_0 \in \bar{\mathcal{D}}$. By possibly redefining \bar{u}^ϵ on a Borel set of measure zero in $\bar{\mathcal{D}}$, we obtain the validity of $(RMV)_\epsilon$ for all $x_0 \in \bar{\mathcal{D}}$, as claimed. □

We conclude this section by estimating the expectation of the accumulated Dirichlet factors Λ_n , in the boundary layer where $d_\epsilon < 1$. This property will be used in Section 7, towards bounding the error of \bar{u}^ϵ from the first order Taylor expansion of the weak solution u to (RL).

Lemma 6.2 Assume (BH). Then for each $\epsilon \ll 1$ and $x_0 \in \bar{\mathcal{D}}$, the function $\sum_{i=0}^\infty \mathbb{1}_{\{d_\epsilon(X_k) < 1\}} \Lambda_k^{x_0}$ is an integrable random variable and we have:

$$\mathbb{E} \left[\sum_{i=0}^\infty \mathbb{1}_{\{d_\epsilon(X_i^{x_0}) < 1\}} \Lambda_i^{x_0} \right] \leq \frac{C}{\epsilon},$$

where C is a constant that is independent of ϵ and x_0 , but may depend on f, \mathcal{D} and γ .

Proof Fix $\epsilon \ll 1$, $x_0 \in \bar{\mathcal{D}}$. Given a constant $\lambda > 0$, we consider the sequence of random variables:

$$M_n = \sum_{i=0}^{n-1} \left(\lambda \mathbb{1}_{\{d_\epsilon(X_i^{x_0}) < 1\}} - s_\epsilon \circ X_{i+1}^{x_0} \right) \Lambda_i,$$

where we adopt the convention that $M_0 = 0$. We claim that $\{M_n\}_{n=0}^\infty$ is a supermartingale with respect to the filtration $\{\mathcal{F}_n\}_{n=0}^\infty$, provided that $\lambda = \lambda(\mathcal{D})$ is chosen appropriately. Indeed:

$$\begin{aligned} \mathbb{E}(M_{n+1} - M_n \mid \mathcal{F}_n) &= \mathbb{E}\left((\lambda \mathbb{1}_{\{d_\epsilon(X_n) < 1\}} - s_\epsilon(X_{n+1})) \Lambda_n \mid \mathcal{F}_n \right) \\ &= \left(\lambda \mathbb{1}_{\{d_\epsilon(X_n) < 1\}} - \frac{1}{\epsilon} \mathbb{E}(s_\epsilon(X_{n+1}) \mid \mathcal{F}_n) \right) \epsilon \Lambda_n \leq 0 \quad \mathbb{P} - \text{a.s. in } \Omega, \end{aligned}$$

where the last inequality follows by observing that on the event $\{\text{dist}(X_n, \partial\mathcal{D}) \geq \epsilon\}$ the quantity in parentheses is clearly nonpositive, whereas on the event $\{\text{dist}(X_n, \partial\mathcal{D}) < \epsilon\}$ it is nonpositive, upon choosing λ small so that:

$$\begin{aligned} \frac{1}{\epsilon} \mathbb{E}(s_\epsilon(X_{n+1}) \mid \mathcal{F}_n) &= \frac{1}{\epsilon} \int_{B_\epsilon(X_n) \cap \mathcal{D}} s_\epsilon(y) \, dy \\ &\geq c \int_{B_\epsilon(X_n) \cap \mathcal{D}} (1 - d_\epsilon(y)^2)^{\frac{N+1}{2}} \, dy \geq \lambda \quad \mathbb{P} - \text{a.s. in } \Omega. \end{aligned}$$

The supermartingale property implies: $\mathbb{E}[M_{n+1}] \leq \mathbb{E}[M_0] = 0$ and thus we get:

$$\begin{aligned} \mathbb{E} \left[\sum_{i=0}^n \epsilon \mathbb{1}_{\{d_\epsilon(X_i) < 1\}} \Lambda_i^{x_0} \right] &\leq \frac{1}{\lambda} \mathbb{E} \left[\sum_{i=0}^n s_\epsilon(X_{i+1}) \Lambda_i \right] \leq \frac{2}{\lambda} \mathbb{E} \left[\sum_{i=0}^n s_\epsilon(X_{i+1}) \Lambda_{i+1} \right] \\ &= \frac{2}{\lambda \gamma} \mathbb{E} \left[\sum_{i=0}^n (1 - (1 - \gamma s_\epsilon(X_{i+1}))) \Lambda_{i+1} \right] \\ &= \frac{2}{\lambda \gamma} \mathbb{E} \left[\sum_{i=0}^n \Lambda_{i+1} - \sum_{i=0}^n \Lambda_{i+2} \right] = \frac{2}{\lambda \gamma} \mathbb{E}[\Lambda_1 - \Lambda_{n+2}] \leq \frac{2}{\lambda \gamma}. \end{aligned}$$

The Lemma follows by passing to the limit with $n \rightarrow \infty$. □

7 Convergence to $W^{2,p}$ Solutions: Case $f \in C(\bar{\mathcal{D}})$

In this section we complete the proof of Theorem 1.1 (ii). The key point is an estimate of the remainder in the Taylor expansion (1.1), for u that is a $W^{2,p}$ solution of (RL). In the boundary layer where $\text{dist}(x_0, \partial\mathcal{D}) < \epsilon$, this quantity is treated by means of Lemma 6.2, whereas in the interior of \mathcal{D} we find a representation of the Newtonian potential component of u via a convolution with suitable probabilistic kernels.

Theorem 7.1 *Assume (BH) and let $u \in C^1(\bar{\mathcal{D}})$ be the unique $W^{2,p}$ solution to (RL). Define the following uniformly bounded sequence of Borel “remainder” functions $\{R_\epsilon\}_{\epsilon \rightarrow 0}$:*

$$R_\epsilon(x) = u(x) - (1 - \gamma s_\epsilon(x)) \int_{B_\epsilon(x) \cap \mathcal{D}} u(y) \, dy - \frac{\epsilon^2}{2(N+2)} f(x).$$

Then, for every $\epsilon \ll 1$ and $x_0 \in \bar{\mathcal{D}}$ we have:

- (i) $|R_\epsilon(x_0)| \leq \epsilon a_\epsilon$, where $\lim_{\epsilon \rightarrow 0} a_\epsilon = 0$, uniformly in $x_0 \in \bar{\mathcal{D}}$.

- (ii) $u(x_0) - \bar{u}^\epsilon(x_0) = \lim_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=0}^n (R_\epsilon \circ X_i^{\epsilon, x_0}) \mathbb{1}_{\{d_\epsilon(X_i^{x_0}) \geq 1\}} \Lambda_i^{\epsilon, x_0} \right].$
- (iii) *There exists a family of positive Borel functions $\{h_\epsilon : B_\epsilon(0) \rightarrow \mathbb{R}\}_{\epsilon \rightarrow 0}$ that are probability densities: $\int_{B_\epsilon(0)} h_\epsilon(y) dy = 1$, and such that whenever $\text{dist}(x_0, \partial \bar{\mathcal{D}}) \geq \epsilon$, there holds:*

$$R_\epsilon(x_0) = \frac{\epsilon^2}{2(N+2)} \left(\int_{B_\epsilon(x_0)} h_\epsilon(x_0 - y) f(y) dy - f(x_0) \right).$$

Proof 1. Clearly, the Taylor expansion: $u(y) = u(x_0) + \langle \nabla u(x_0), y - x \rangle + o(\epsilon)$ holds uniformly in $y \in B_\epsilon(x_0) \cap \bar{\mathcal{D}}$ and $x_0 \in \bar{\mathcal{D}}$. Arguing as in the proof of Theorem 3.1 it follows that:

$$\begin{aligned} \int_{B_\epsilon(x_0) \cap \bar{\mathcal{D}}} u(y) dy &= u(x_0) + \langle \nabla u(x_0), \int_{B_\epsilon(x_0) \cap \bar{\mathcal{D}}} y - x_0 dy \rangle + o(\epsilon) \\ &= u(x_0) + \gamma s_\epsilon(x) u(x_0) + O(\epsilon s_\epsilon(x_0)) + o(\epsilon). \end{aligned}$$

Consequently, we obtain the bound in (i):

$$R_\epsilon(x_0) = u(x_0) - (1 - \gamma^2 s_\epsilon(x_0)^2) u(x_0) + O(\epsilon s_\epsilon(x_0)) + o(\epsilon) = o(\epsilon).$$

2. To show (ii), observe that the following sequence of random variables $\{M_n\}_{n=0}^\infty$ is a martingale with respect to the filtration $\{\mathcal{F}_n\}_{n=0}^\infty$:

$$M_n = (u \circ X_n) \Lambda_n + \frac{\epsilon^2}{2(N+2)} \sum_{i=0}^{n-1} (f \circ X_i) \Lambda_i + \sum_{i=0}^{n-1} (R_\epsilon \circ X_i) \Lambda_i,$$

where we adopt the convention that $M_0 = 0$. Indeed:

$$\begin{aligned} \mathbb{E}(M_{n+1} - M_n \mid \mathcal{F}_n) &= \mathbb{E}(u \circ X_{n+1} \mid \mathcal{F}_n) \Lambda_{n+1} - (u \circ X_n) \Lambda_n + \frac{\epsilon^2}{2(N+2)} (f \circ X_n) \Lambda_n + (R_\epsilon \circ X_n) \Lambda_n \\ &= \left((1 - \gamma s_\epsilon(X_n)) \int_{B_\epsilon(X_n) \cap \bar{\mathcal{D}}} u(y) dy - u \circ X_n + \frac{\epsilon^2}{2(N+2)} (f \circ X_n) + R_\epsilon \right) \Lambda_n \\ &= 0 \quad \mathbb{P} - \text{a.s. in } \Omega. \end{aligned}$$

The above yields $u(x_0) = \mathbb{E}[M_0] = \mathbb{E}[M_{n+1}]$, resulting in:

$$u(x_0) - \frac{\epsilon^2}{2(N+2)} \mathbb{E} \left[\sum_{i=0}^n (f \circ X_i^{\epsilon, x_0}) \Lambda_i^{\epsilon, x_0} \right] = \mathbb{E} \left[(u \circ X_{n+1}^{\epsilon, x_0}) \Lambda_{n+1}^{\epsilon, x_0} \right] + \mathbb{E} \left[\sum_{i=0}^n (R_\epsilon \circ X_i) \Lambda_i \right].$$

Passing to the limit with $n \rightarrow \infty$ and using (6.3) and (6.4) implies:

$$\begin{aligned} u(x_0) - \bar{u}^\epsilon(x_0) &= \lim_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=0}^n (R_\epsilon \circ X_i^{\epsilon, x_0}) \Lambda_i \right] \\ &= \lim_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=0}^n O(\epsilon a_\epsilon) \mathbb{1}_{\{d_\epsilon(X_i) < 1\}} \Lambda_i \right] + \lim_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=0}^n (R_\epsilon \circ X_i) \mathbb{1}_{\{d_\epsilon(X_i) \geq 1\}} \Lambda_i \right] \end{aligned}$$

and we conclude (ii) by Lemma 6.2.

3. Recall that the Newtonian potential $\Gamma * f \in \mathcal{C}^1(\mathbb{R}^N)$, namely $(\Gamma * f)(x) = \int_{\mathcal{D}} \Gamma(x - y) f(y) dy$, is given in terms of the fundamental solution:

$$\Gamma(x) = \begin{cases} \frac{1}{N(N-2)|B_1^N|} |x|^{2-N} & \text{when } N > 2 \\ -\frac{1}{2\pi} \log |x| & \text{when } N = 2, \end{cases}$$

and it satisfies: $-\Delta(\Gamma * f) = f$ in \mathcal{D} , in the sense of distributions [11, section 4.1]. Writing:

$$u = (\Gamma * f)|_{\bar{\mathcal{D}}} + w,$$

where $w \in C^1(\bar{\mathcal{D}})$ is harmonic in \mathcal{D} , we derive the below formula, valid for $x_0 \in \mathcal{D}$ with $\text{dist}(x_0, \partial\mathcal{D}) \geq \epsilon$, in virtue of the mean value property of w and by changing the integration order in convolution:

$$\begin{aligned} u(x_0) - \int_{B_\epsilon(x_0) \cap \mathcal{D}} u(y) \, dy &= (\Gamma * f)(x_0) - \int_{B_\epsilon(x_0)} (\Gamma * f)(y) \, dy \\ &= \int_{\mathcal{D}} \left(\Gamma(x_0 - y) - \int_{B_\epsilon(x_0 - y)} \Gamma(z) \, dz \right) f(y) \, dy \\ &= \frac{\epsilon^2}{2(N + 2)} \int_{\mathcal{D}} h_\epsilon(x_0 - y) f(y) \, dy. \end{aligned}$$

For the final convolution kernel, we have denoted:

$$h_\epsilon(x) = \frac{2(N + 2)}{\epsilon^2} \left(\Gamma(x) - \int_{B_\epsilon(x)} \Gamma(y) \, dy \right).$$

The assertion (iii) follows provided we show the claimed properties of the sequence $\{h_\epsilon\}_{\epsilon \rightarrow 0}$. When $|x| \geq \epsilon$, then Γ is harmonic on $B_\epsilon(x)$, so that $h_\epsilon(x) = 0$ by the mean value property. On the other hand, for $|x| < \epsilon$, we get:

$$\begin{aligned} \int_{B_\epsilon(x)} \Gamma(y) \, dy &= \int_{B_{|x|}(0)} \Gamma(x - z) \, dz + \int_{|x|}^\epsilon \int_{\partial B_r(0)} \Gamma(x - z) \, d\sigma(z) \, dr \tag{7.1} \\ &= |B_1^N| \cdot |x|^N \Gamma(x) + \int_{|x|}^\epsilon |\partial B_1^N| \cdot r^{N-1} \int_{\partial B_r(0)} \Gamma(y) \, d\sigma(y) \, dr. \end{aligned}$$

The first term in Eq. 7.1 results by the mean value property of the harmonic function Γ on $B_{|x|}(x)$. The second term follows by noting that $g(x) = \int_{\partial B_r(0)} \Gamma(x - y) \, d\sigma(y)$ is constant on $B_r(0)$. This fact is the Newton shell theorem, that can be proved as follows. The function g is clearly harmonic on $B_r(0)$ and it is also radially symmetric: $g(x) = g(|x|)$ by the radial symmetry of Γ . Hence, the spherical mean value property yields: $g(0) = \int_{B_s(0)} g(y) \, dy = g(y_0)$ for all $|y_0| = s < r$.

In conclusion, we obtain:

$$\int_{B_\epsilon(x)} \Gamma(y) \, dy = \begin{cases} \frac{1}{N(N-2)} \left(|x|^2 + \int_{|x|}^\epsilon Nr \, dr \right) = \frac{\epsilon^2}{2(N-2)} - \frac{|x|^2}{2N} & \text{when } N > 2 \\ -\frac{1}{2} \left(|x|^2 \log|x| + \int_{|x|}^\epsilon 2r \log|r| \, dr \right) = \frac{1}{2} \left(\frac{\epsilon^2}{2} - \epsilon^2 \log|\epsilon| - \frac{|x|^2}{2} \right) & \text{when } N = 2, \end{cases}$$

and further:

$$h_\epsilon(x) = \begin{cases} \frac{2(N+2)}{N(N-2)|B_1^N|} \left(\frac{|x|^{2-N}}{\epsilon^2} - \frac{N}{2\epsilon^N} + \frac{(N-2)|x|^2}{2\epsilon^{N-2}} \right) & \text{when } N > 2 \\ \frac{4}{\pi} \left(\frac{\log|\epsilon|}{\epsilon^2} + \frac{|x|^2}{2\epsilon^4} - \frac{1}{2\epsilon^2} - \frac{\log|x|}{\epsilon^2} \right) & \text{when } N = 2. \end{cases}$$

It follows by a calculation that $h_\epsilon(x) > 0$ for all $x \in B_\epsilon(0)$ and $\int_{B_\epsilon} h_\epsilon = 1$. □

We are now ready to prove the remaining part of Theorem 1.1.

Corollary 7.2 Assume (BH) and let $f \in C(\bar{\mathcal{D}})$. Then $\{\bar{u}^\epsilon\}_{\epsilon \rightarrow 0}$ converge uniformly on $\bar{\mathcal{D}}$ to u that is the unique $W^{2,p}$ solution of (RL).

Proof By Theorem 7.1 (iii) we get, for all $x_0 \in \mathcal{D}$ with $\text{dist}(x_0, \partial\mathcal{D}) \geq \epsilon$:

$$|R_\epsilon(x_0)| = o(\epsilon^2),$$

where the Landau symbol o is uniform in x_0 but may depend on f and N . Further:

$$|u(x_0) - \bar{u}_\epsilon(x_0)| \leq \lim_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=0}^n |R_\epsilon \circ X_i^{\epsilon, x_0}| \mathbb{1}_{\{d_\epsilon(X_i^{x_0}) \geq 1\}} \Lambda_i^{\epsilon, x_0} \right] = o(1),$$

in view of Theorem 7.1 (ii) and of Eq. 6.3. The proof is done. □

8 The Lower Bound

In this section, we prove Theorem 1.2 and the optimality of the inverse quadratic estimate in Corollary 5.2, through a precise lower bound on the stopping time τ^{ϵ, x_0} . As a consequence, we obtain a uniform bound for solutions u_ϵ to (RMV) $_\epsilon$ with $f \geq 0$, yielding the lower bound for solutions u of the Robin problem (RL). The same optimal bound will be deduced directly for u via analytical arguments in Lemma 8.4.

Definition 8.1 Let \mathcal{D} be as in (BH) and fix a radius $\rho > 0$ that is strictly smaller than some uniform inner supporting sphere radius r of \mathcal{D} , given by the property:

$$\text{for every } x \in \partial\mathcal{D} \text{ exists } B_r(a) \subset \mathcal{D} \text{ such that } |x - a| = r. \tag{8.1}$$

We define a continuous function $Z^\rho : \bar{\mathcal{D}} \rightarrow \mathcal{D}$ satisfying:

$$x \in \bar{B}_\rho(Z^\rho(x)) \subset \bar{\mathcal{D}} \quad \text{for all } x \in \bar{\mathcal{D}},$$

as follows. For every $x \in \bar{\mathcal{D}}$ such that $\text{dist}(x, \partial\mathcal{D}) \leq \rho$, there exists exactly one $Z^\rho(x) \in \mathcal{D}$ that is the center of the inner ρ -supporting sphere at $\pi_{\partial\mathcal{D}}x$. For $x \in \mathcal{D}$ with $\text{dist}(x, \partial\mathcal{D}) > \rho$, we set $Z^\rho(x) = x$. It is straightforward that for every $\epsilon \ll 1, x_0 \in \bar{\mathcal{D}}$ and $n \geq 0$ there holds:

$$|Z^\rho(X_{n+1}^{\epsilon, x_0}) - X_{n+1}^{\epsilon, x_0}| \leq |Z^\rho(X_n^{\epsilon, x_0}) - X_{n+1}^{\epsilon, x_0}|. \tag{8.2}$$

Theorem 8.2 Assume (BH) and let $\bar{r} < r$ with r as in Eq. 8.1. Then for every $\epsilon \ll 1$ there holds:

$$\mathbb{E}[\tau^{\epsilon, x_0}] \geq \frac{2(N+2)}{\gamma N} \cdot \frac{\bar{r}}{\epsilon^2} \quad \text{for all } x_0 \in \bar{\mathcal{D}}. \tag{8.3}$$

Proof 1. Fix an intermediate radius $r_* \in (\bar{r}, r)$ and consider the auxiliary sequence of random variables $\{Z_n^{\epsilon, r_*, x_0} = Z^{r_*} \circ X_n^{\epsilon, x_0}\}_{n=0}^\infty$ given in Definition 8.1 (see also Fig. 5). As




Fig. 5 The auxiliary balls $B_{r_*}(Z_n)$ and the process $\{X_n\}_{n=0}^\infty$ in Theorem 8.2

usual, we will drop the superscripts ϵ, r_* and x_0 to alleviate the notation. We now define the following sequence:

$$M_n = |\mathcal{Z}_n - X_n|^2 - n \frac{N\epsilon^2}{N+2} + \frac{2\bar{r}}{\gamma} \cdot \sum_{j=1}^n \mathbb{1}_{b_j < \gamma s_\epsilon(X_{j-1})}.$$

and aim to prove that $\{M_n\}_{n=0}^\infty$ is a supermartingale with respect to the filtration $\{\mathcal{F}_n\}_{n=0}^\infty$. Firstly, because of Eq. 8.2 it follows that:

$$\begin{aligned} \mathbb{E}(M_{n+1} - M_n \mid \mathcal{F}_n) &= \mathbb{E}(|\mathcal{Z}_{n+1} - X_{n+1}|^2 \mid \mathcal{F}_n) - |\mathcal{Z}_n - X_n|^2 \\ &\quad - \frac{N\epsilon^2}{N+2} + \frac{2\bar{r}}{\gamma} \cdot \mathbb{E}(\mathbb{1}_{b_{n+1} < \gamma s_\epsilon(X_n)} \mid \mathcal{F}_n) \\ &\leq \mathbb{E}(|\mathcal{Z}_n - X_{n+1}|^2 \mid \mathcal{F}_n) - |\mathcal{Z}_n - X_n|^2 - \frac{N\epsilon^2}{N+2} + 2\bar{r} \cdot s_\epsilon(X_n). \end{aligned} \tag{8.4}$$

Then, with the help of Eqs. 2.3, 2.5 and 5.1 we get:

$$\begin{aligned} &\mathbb{E}(|\mathcal{Z}_n - X_{n+1}|^2 \mid \mathcal{F}_n) - |\mathcal{Z}_n - X_n|^2 \\ &= \int_{B_\epsilon(0)} |\mathcal{Z}_n - (X_n + \epsilon w_n^{X_{n-1}})|^2 dw_{n+1} - |\mathcal{Z}_n - X_n^{x_0}|^2 \\ &= \int_{B_\epsilon(X_n^{x_0}) \cap \mathcal{D}} |y - \mathcal{Z}_n|^2 dy - |\mathcal{Z}_n - X_n|^2 \\ &= \int_{B_\epsilon(X_n) \cap \mathcal{D}} |y - X_n|^2 dy + 2 \left\langle \int_{B_\epsilon(X_n) \cap \mathcal{D}} y - X_n dy, X_n - \mathcal{Z}_n \right\rangle \\ &= \frac{N\epsilon^2}{N+2} - 2s_\epsilon(X_n) \langle \bar{n}(\pi_{\partial\mathcal{D}} X_n), X_n - \mathcal{Z}_n \rangle + \mathcal{O}(\epsilon s_\epsilon(X_n)). \end{aligned}$$

Indeed, from Eq. 2.5 we directly obtain:

$$\int_{B_\epsilon(X_n) \cap \mathcal{D}} |y - X_n|^2 dy = \text{trace} \int_{B_\epsilon(X_n) \cap \mathcal{D}} (y - X_n)^{\otimes 2} dy = \frac{N\epsilon^2}{N+2} + \mathcal{O}(\epsilon s_\epsilon(X_n)),$$

while (2.3) yields:

$$\left\langle \int_{B_\epsilon(X_n) \cap \mathcal{D}} y - X_n dy, X_n - \mathcal{Z}_n \right\rangle = -s_\epsilon(X_n) \langle \bar{n}(\pi_{\partial\mathcal{D}} X_n), X_n - \mathcal{Z}_n \rangle + \mathcal{O}(\epsilon s_\epsilon(X_n)) |X_n - \mathcal{Z}_n|.$$

Since $X_n - \mathcal{Z}_n = |X_n - \mathcal{Z}_n| \cdot \bar{n}(\pi_{\partial\mathcal{D}} X_n)$, and since $|X_n - \mathcal{Z}_n| \geq r_* - \epsilon$ when $s_\epsilon(X_n) \neq 0$, it follows that for all $\epsilon \ll r_* - \bar{r}$ we get:

$$\begin{aligned} &\mathbb{E}(|\mathcal{Z}_n - X_{n+1}|^2 \mid \mathcal{F}_n) - |\mathcal{Z}_n - X_n|^2 \frac{N\epsilon^2}{N+2} - 2s_\epsilon(X_n) \cdot |X_n - \mathcal{Z}_n| \\ &\quad + \mathcal{O}(\epsilon s_\epsilon(X_n)) \leq \frac{N\epsilon^2}{N+2} - 2r_* s_\epsilon(X_n) + \mathcal{O}(\epsilon s_\epsilon(X_n)) \leq \frac{N\epsilon^2}{N+2} - 2\bar{r} s_\epsilon(X_n). \end{aligned}$$

Recalling (8.4), we get the desired supermartingale property: $\mathbb{E}(M_{n+1} - M_n \mid \mathcal{F}_n) \leq 0$.

2. For each fixed $k \geq 1$, apply now Doob's Optional Stopping to $\{M_n\}_{n=0}^\infty$ and the finite stopping time $\tau^{\epsilon, x_0} \wedge k$, to the effect that:

$$\begin{aligned} r_*^2 &\geq |\mathcal{Z}_0 - x_0|^2 = \mathbb{E}[M_0] \geq \mathbb{E}[M_{\tau^{\epsilon, x_0} \wedge k}] \\ &= \mathbb{E} \left[\left| \mathcal{Z}_{\tau^{\epsilon, x_0} \wedge k} - X_{\tau^{\epsilon, x_0} \wedge k} \right|^2 \right] - \mathbb{E} \left[\tau^{\epsilon, x_0} \wedge k \right] \frac{N\epsilon^2}{N+2} + \frac{2\bar{r}}{\gamma} \cdot \mathbb{P}(\tau^{\epsilon, x_0} \leq k). \end{aligned}$$

Passing to the limit with $k \rightarrow \infty$, we obtain:

$$r_*^2 \geq \mathbb{E}[|Z_{\tau^{\epsilon, x_0}} - X_{\tau^{\epsilon, x_0}}|^2] - \mathbb{E}[\tau^{\epsilon, x_0}] \frac{N\epsilon^2}{N+2} + \frac{2\bar{r}}{\gamma} \geq |r_* - \epsilon|^2 - \mathbb{E}[\tau^{\epsilon, x_0}] \frac{N\epsilon^2}{N+2} + \frac{2\bar{r}}{\gamma}.$$

Writing $|r_* - \epsilon|^2 \geq r_*^2 - 2r_*\epsilon$, it follows that:

$$\mathbb{E}[\tau^{\epsilon, x_0}] \cdot \frac{N\epsilon^2}{N+2} \geq \frac{2\bar{r}}{\gamma} - 2r_*\epsilon,$$

which implies (8.3) for \bar{r} replaced by any smaller radius, provided that $\epsilon \ll 1$. This completes the proof of the claim, for any $\bar{r} < r$. □

Corollary 8.3 Assume (BH), let $f \geq 0$ and let $r > 0$ satisfy (8.1). Then:

(i) The solution to (RMV) $_{\epsilon}$ satisfies:

$$u_{\epsilon}(x_0) \geq \frac{\bar{r}}{\gamma N} \cdot \inf_{\bar{D}} f \quad \text{for all } x_0 \in \bar{D},$$

for any radius $\bar{r} < r$, provided that $\epsilon \ll 1$.

(ii) If some sequence of solutions $\{u_{\epsilon}\}_{\epsilon \rightarrow 0}$ to (RMV) $_{\epsilon}$ converges pointwise on \bar{D} , then the limit $u : \bar{D} \rightarrow \mathbb{R}$ satisfies:

$$u(x_0) \geq \frac{r}{\gamma N} \cdot \inf_{\bar{D}} f \quad \text{for all } x_0 \in \bar{D}. \tag{8.5}$$

Proof From Eq. 8.3 and the definition 5.4 it directly follows that:

$$u^{\epsilon}(x_0) \geq \frac{\epsilon^2}{2(N+2)} \mathbb{E}[\tau^{\epsilon, x_0} - 1] \cdot \inf_{\bar{D}} f \geq \left(\frac{\bar{r}}{\gamma N} - \frac{\epsilon^2}{2(N+2)} \right) \cdot \inf_{\bar{D}} f,$$

resulting in (i) for \bar{r} replaced by any smaller radius, for $\epsilon \ll 1$. Recalling Lemma 5.1, this completes the proof of the claim for any $\bar{r} < r$. The limiting statement (ii) is self-evident from (i). □

Automatically, the limiting solution u of $\{u^{\epsilon}\}_{\epsilon \rightarrow 0}$ obeys the bound in Eq. 8.5. We now present an analytical proof of the same result, under stronger regularity assumptions. It is based on the maximum principle (see, for example, [11, Corollary 3.2]).

Lemma 8.4 Let $\mathcal{D} \subset \mathbb{R}^N$ be an open, bounded and connected set, satisfying the uniform inner supporting sphere condition with radius $r > 0$ as in Eq. 8.1. Given two constants: $\gamma > 0$ and $m \geq 0$, assume that $u \in C^1(\bar{\mathcal{D}})$ satisfies (in the sense of distributions in \mathcal{D}):

$$-\Delta u \geq m \quad \text{in } \mathcal{D}, \quad \frac{\partial u}{\partial \bar{n}} + \gamma u = 0 \quad \text{on } \partial \mathcal{D}.$$

Then: $\min_{\bar{\mathcal{D}}} u \geq \frac{rm}{\gamma N}$.

Proof By the maximum principle, there holds: $\min_{\bar{\mathcal{D}}} u = \min_{\partial \mathcal{D}} u = u(x_0)$ for some $x_0 \in \partial \mathcal{D}$. Consider the inner supporting ball $B = B(x_0 - r\bar{n}(x_0))$ and the function $v_r \in C^2(\bar{B})$ given by:

$$v_r(x_0 - r\bar{n}(x_0) + y) = v(|y|) \quad \text{where: } v(s) = \frac{m}{2N}(r^2 - s^2) + \frac{rm}{\gamma N}.$$

Since:

$$-\Delta v_r = m \quad \text{in } B, \quad \frac{\partial v_r}{\partial \vec{n}} + \gamma v_r = 0 \quad \text{on } \partial B,$$

it follows that: $-\Delta(u - v_r) \geq 0$. Applying again the maximum principle, we get:

$$\min_{\bar{B}}(u - v_r) = \min_{\partial B}(u - v_r) = \min_{\partial B} u - \frac{rm}{\gamma N} = u(x_0) - \frac{rm}{\gamma N}.$$

In particular, the difference $u - v_r$ is minimized at x_0 , so: $\frac{\partial(u-v_r)}{\partial \vec{n}}(x_0) \leq 0$, and consequently:

$$(u - v_r)(x_0) = -\frac{1}{\gamma} \cdot \frac{\partial(u - v_r)}{\partial \vec{n}}(x_0) \geq 0.$$

In conclusion: $u(x_0) \geq v(x_0) = \frac{rm}{\gamma N}$, proving the claim. □

Remark 8.5 The bound in Lemma 8.4 is optimal. Take $\mathcal{D} = B_r(0)$ and $u(x) = a - |x|^2$. Then:

$$-\Delta u = 2N \quad \text{in } \mathcal{D}, \quad \frac{\partial u}{\partial \vec{n}} + \gamma u = -2r + \gamma(a - r^2) \quad \text{on } \partial \mathcal{D},$$

so (RL) holds with $f = 2N$ and $\gamma = \frac{2r}{a-r^2} > 0$ by taking $a > r^2$. Then, clearly:

$$\min_{\bar{\mathcal{D}}} u = a - r^2 = \frac{r}{2rN} (a - r^2) 2N = \frac{r}{\gamma N} \min_{\bar{\mathcal{D}}} f.$$

We also remark that using the arguments in [5, Lemma 3.2], [20], one can prove the lower bound on u involving the integral $\int_{\mathcal{D}} f(y) \, dy$, rather than the pointwise quantity $\min_{\bar{\mathcal{D}}} f$.

Appendix : A Proof of Uniqueness of Viscosity Solutions to (RL)

We first make an observation that relies on the assumed regularity of $\partial \mathcal{D}$.

Lemma A.1 *When $\mathcal{D} \subset \mathbb{R}^N$ satisfies the uniform outer supporting sphere condition:*

$$\text{for every } x \in \partial \mathcal{D} \text{ exists } B_r(b) \subset \mathbb{R}^n \setminus \bar{\mathcal{D}} \text{ such that } |x - b| = r, \quad (9.1)$$

with some radius $r > 0$, then the boundary requirements in Definition 4.1 (i) and (ii) can be reduced to: $\langle p, \vec{n}(x) \rangle + \gamma u(x) \leq 0$ and $\langle p, \vec{n}(x) \rangle + \gamma u(x) \geq 0$, respectively.

Proof Let $x \in \partial \mathcal{D}$ and $(p, X) \in J_{\bar{\mathcal{D}}}^{2,+} u(x)$. For each large j , consider the jet:

$$(p_j, X_j) = \left(p - \frac{1}{j} \vec{n}(x), X + r Id_N - (r + j) \vec{n}(x) \otimes^2 \right).$$

We claim that $(p_j, X_j) \in J_{\bar{\mathcal{D}}}^{2,+} u(x)$. In this case, we have:

$$-\text{trace} X_j = -(\text{trace} X + (N - 1)r - j) > f(x),$$

so by (i) there must be: $0 \geq \langle p_j, \vec{n}(x) \rangle + \gamma u(x) = \langle p, \vec{n}(x) \rangle + \gamma u(x) - \frac{1}{j}$. Passing to the limit with $j \rightarrow \infty$ we get the claimed boundary condition. To show that (p_j, X_j) is indeed a super-jet, let:

$$\begin{aligned} q_j(y - x) &= \left\langle -\frac{1}{j} \vec{n}(x), y - x \right\rangle + \frac{1}{2} \left(r Id_N - (r + j) \vec{n}(x) \otimes^2 : (y - x) \otimes^2 \right) \\ &= -\frac{1}{j} \langle \vec{n}(x), y - x \rangle + \frac{r}{2} |(y - x)_{tan}|^2 - \frac{j}{2} \langle \vec{n}(x), y - x \rangle^2. \end{aligned}$$

For $y \in \bar{\mathcal{D}}$ satisfying $\langle \vec{n}(x), y - x \rangle \leq 0$, we get: $q_j(y - x) \geq |\langle \vec{n}(x), y - x \rangle| \left(\frac{1}{j} - \frac{j}{2} |\langle \vec{n}(x), y - x \rangle| \right) \geq 0$, if $|y - x|$ is small enough. On the other other hand, for $y \in \bar{\mathcal{D}}$ such that $\langle \vec{n}(x), y - x \rangle \geq 0$, we get:

$$\begin{aligned} q_j(y - x) &\geq -\frac{2}{j} \langle \vec{n}(x), y - x \rangle + \frac{r}{2} |(y - x)_{tan}|^2 \\ &\geq -\frac{2}{j} \left(r - \sqrt{r^2 - |(y - x)_{tan}|^2} \right) + \frac{r}{2} |(y - x)_{tan}|^2 \geq 0, \end{aligned}$$

as for small $|y - x|$ and $j \gg 1$ there holds: $r - \sqrt{r^2 - |(y - x)_{tan}|^2} \leq \frac{1}{r} |(y - x)_{tan}|^2 \leq \frac{jr}{4} |(y - x)_{tan}|^2$. Thus, in both cases, the validity of Eq. 4.1 for (p, X) , implies the same asymptotic bound for each (p_j, X_j) with sufficiently large j . □

Lemma A.2 *Assume the uniform outer supporting sphere condition (9.1) with radius $r > 0$. Then the Robin problem (RL) with $f \in \mathcal{C}(\bar{\mathcal{D}})$ has at most one viscosity solution.*

Proof 1. Let u, v be two viscosity solutions to (RL). We will prove that $u \leq v$. In fact, the same analysis works when u is a viscosity sub-solution and v is a viscosity super-solution, in the sense of Definition 4.1 (i) and (ii), where u is assumed only to be upper-semicontinuous and v lower-semicontinuous, and where the jets sets $J_{\bar{\mathcal{D}}}^{2,+}$ and $J_{\bar{\mathcal{D}}}^{2,-}$ are replaced by their closures $\bar{J}_{\bar{\mathcal{D}}}^{2,+}$ and $\bar{J}_{\bar{\mathcal{D}}}^{2,-}$, respectively (see [9] for the details). Also, we recall that the requirements at boundary points in Definition 4.1 can be reduced as in Lemma 9.1, because of Eq. 9.1. The stated comparison principle is proved in three steps: by replacing u and v with strict sub- and super-solutions u_δ and v_δ , and by doubling the variables technique with an appropriate nonlinear corrector, separately in the cases when u_δ and v_δ achieves its maximum in \mathcal{D} or on $\partial\mathcal{D}$. We now sketch these arguments.

For a sufficiently large $C \gg 1$ and each $\delta \ll 1$, define:

$$u_\delta(x) = u(x) - \frac{\delta}{2} (|x|^2 - C), \quad v_\delta(x) = v(x) + \frac{\delta}{2} (|x|^2 - C).$$

Assume that $(p, X) \in J_{\bar{\mathcal{D}}}^{2,+} u_\delta(x)$, which is equivalent to: $(p - \delta x, X - \delta Id_N) \in J_{\bar{\mathcal{D}}}^{2,+} u(x)$. Then:

$$\begin{aligned} -\text{trace } X &\leq f(x) - N\delta && \text{when } x \in \mathcal{D} \\ \langle p, \vec{n}(x) \rangle + \gamma u_\delta(x) &\leq -\delta && \text{when } x \in \partial\mathcal{D}. \end{aligned} \tag{9.2}$$

Thus, each u_δ is a strict sub-solution of (RL). Similarly, each v_δ is a strict super-solution, namely $(p, X) \in J_{\bar{\mathcal{D}}}^{2,-} v_\delta(x)$ implies:

$$\begin{aligned} -\text{trace } X &\geq f(x) + N\delta && \text{when } x \in \mathcal{D} \\ \langle p, \vec{n}(x) \rangle + \gamma v_\delta(x) &\geq \delta && \text{when } x \in \partial\mathcal{D}. \end{aligned} \tag{9.3}$$

2. We will show that $u_\delta \leq v_\delta$ in $\bar{\mathcal{D}}$, for all $\delta \ll 1$. By contradiction, fix $\delta > 0$ and assume that:

$$\max_{\bar{\mathcal{D}}} (u_\delta - v_\delta) > 0.$$

We first treat the case of $\max_{\bar{\mathcal{D}}} (u_\delta - v_\delta) > \max_{\partial\bar{\mathcal{D}}} (u_\delta - v_\delta)$. We apply [9, Proposition 3.7] to: $\Phi(x, y) = u_\delta(x) - v_\delta(y)$ and $\Psi(x, y) = \frac{1}{2}|x - y|^2$ and obtain a sequence $\{(x_\alpha, y_\alpha)\}_{\alpha \rightarrow \infty}$ of maximizers to $\Phi - \alpha\Psi$ on $\bar{\mathcal{D}} \times \bar{\mathcal{D}}$ that converges to some diagonal element (z_0, z_0)

such that z_0 is a maximizer of $u_\delta - v_\delta$ and thus $z_0 \in \mathcal{D}$. Applying [9, Theorem 3.2] to: $w(x, y) = u_\delta(x) - v_\delta(y)$ and $\phi(x, y) = \frac{\alpha}{2}|x - y|^2$, we further obtain sequences of matrices $X_\alpha, Y_\alpha \in \mathbb{R}_{\text{sym}}^{N \times N}$ satisfying:

$$\begin{aligned} (\alpha(x_\alpha - y_\alpha), X_\alpha) &\in J_{\mathcal{D}}^{2,+} u_\delta(x_\alpha), & (\alpha(x_\alpha - y_\alpha), Y_\alpha) &\in J_{\mathcal{D}}^{2,-} v_\delta(y_\alpha) \\ \begin{bmatrix} X_\alpha & 0 \\ 0 & -Y_\alpha \end{bmatrix} &\leq 3\alpha \begin{bmatrix} Id_N & -Id_N \\ -Id_N & Id_N \end{bmatrix}. \end{aligned} \tag{9.4}$$

The first two assertions above, together with (9.2), (9.3) yield:

$$-\text{trace}(X_\alpha - Y_\alpha) \leq f(x_\alpha) - f(y_\alpha) - 2N\delta \rightarrow -2\delta \quad \text{as } \alpha \rightarrow \infty,$$

which contradicts the last assertion in Eq. 9.4 that implies: $\text{trace}(X_\alpha - Y_\alpha) \leq 0$.

3. We now treat the remaining case, namely that of:

$$\max_{\mathcal{D}} (u_\delta - v_\delta) = (u_\delta - v_\delta)(z_0) > 0 \quad \text{for some } z_0 \in \partial\mathcal{D}. \tag{9.5}$$

Applying [9, Proposition 3.7] to:

$$\Phi(x, y) = u_\delta(x) - v_\delta(y) - \gamma u_\delta(z_0)\langle y - x, \vec{n}(z_0) \rangle - |x - z_0|^2, \quad \Psi(x, y) = \frac{1}{2}|x - y|^2,$$

we obtain a sequence $\{(x_\alpha, y_\alpha)\}_{\alpha \rightarrow \infty}$ of maximizers to $\Phi - \alpha\Psi$ that converges to some (z, z) , where z is a maximizer of $\Phi(z, z) = u_\delta(z) - v_\delta(z) - |z - z_0|^4$. Hence there must be $z = z_0$. Also:

$$\alpha|x_\alpha - y_\alpha|^2 \rightarrow 0 \quad \text{as } \alpha \rightarrow \infty. \tag{9.6}$$

We now apply [9, Theorem 3.2] to:

$$w(x, y) = u_\delta(x) - v_\delta(y), \quad \phi(x, y) = \frac{\alpha}{2}|x - y|^2 + \gamma u_\delta(z_0)\langle y - x, \vec{n}(z_0) \rangle,$$

which yields existence of sequences of matrices $X_\alpha, Y_\alpha \in \mathbb{R}_{\text{sym}}^{N \times N}$ satisfying:

$$\begin{aligned} (\alpha(x_\alpha - y_\alpha) - \gamma u_\delta(z_0)\vec{n}(z_0) + 4|x_\alpha - z_0|^2(x_\alpha - z_0), X_\alpha) &\in J_{\mathcal{D}}^{2,+} u_\delta(x_\alpha), \\ (\alpha(x_\alpha - y_\alpha) - \gamma u_\delta(z_0)\vec{n}(z_0), Y_\alpha) &\in J_{\mathcal{D}}^{2,-} v_\delta(y_\alpha), \\ \begin{bmatrix} X_\alpha & 0 \\ 0 & -Y_\alpha \end{bmatrix} &\leq \nabla^2 \phi(x_\alpha, y_\alpha) + \frac{1}{\alpha} \nabla^2 \phi(x_\alpha, y_\alpha)^2. \end{aligned} \tag{9.7}$$

Since $\nabla^2 \phi(x_\alpha, y_\alpha) = \alpha \begin{bmatrix} Id_N & -Id_N \\ -Id_N & Id_N \end{bmatrix} + \mathcal{O}(|x_\alpha - z_0|^2)$, the last assertion above implies:

$$\text{trace}(X_\alpha - Y_\alpha) \leq \mathcal{O}(|x_\alpha - z_0|^2 + \frac{1}{\alpha}|x_\alpha - z_0|^4) \rightarrow 0 \quad \text{as } \alpha \rightarrow \infty. \tag{9.8}$$

Note that for large $\alpha \gg 1$ there must be $x_\alpha, y_\alpha \in \mathcal{D}$. Indeed, if $x_\alpha \in \partial\mathcal{D}$ then Eq. 9.2 and the first assertion in Eq. 9.7 yield the contradiction in:

$$\begin{aligned} -\delta &\geq \alpha \langle x_\alpha - y_\alpha, \vec{n}(x_\alpha) \rangle - \gamma u_\delta(z_0)\langle \vec{n}(z_0), \vec{n}(x_\alpha) \rangle \\ &\quad + 4|x_\alpha - z_0|^2 \langle x_\alpha - z_0, \vec{n}(x_\alpha) \rangle + \gamma u_\delta(x_\alpha) \geq -\frac{\alpha}{2r}|x_\alpha - y_\alpha|^2 \\ &\quad + \gamma u_\delta(z_0)\langle \vec{n}(z_0), \vec{n}(x_\alpha) \rangle + \mathcal{O}(|x_\alpha - z_0|^3) + \gamma u_\delta(x_\alpha) \\ &\rightarrow 0 \quad \text{as } \alpha \rightarrow \infty, \end{aligned}$$

where we used (9.1) for the bound $\langle y_\alpha - x_\alpha, \vec{n}(x_\alpha) \rangle \leq \frac{1}{2r} |x_\alpha - y_\alpha|^2$, followed by Eq. 9.8. Similarly, if $y_\alpha \in \partial\mathcal{D}$ then Eq. 9.3 and the second assertion in Eq. 9.7 brings the contradiction with Eq. 9.5, as:

$$\begin{aligned} \delta &\leq \alpha \langle x_\alpha - y_\alpha, \vec{n}(y_\alpha) \rangle - \gamma u_\delta(z_0) \langle \vec{n}(z_0), \vec{n}(y_\alpha) \rangle + \gamma v_\delta(y_\alpha) \\ &\leq \frac{\alpha}{2r} |x_\alpha - y_\alpha|^2 - \gamma u_\delta(z_0) \langle \vec{n}(z_0), \vec{n}(y_\alpha) \rangle + \gamma v_\delta(y_\alpha) \\ &\rightarrow \gamma (v_\delta(z_0) - u_\delta(z_0)) \quad \text{as } \alpha \rightarrow \infty. \end{aligned}$$

The fact of $x_\alpha, y_\alpha \in \mathcal{D}$ established, we use Eq. 9.7 together with Eqs. 9.2, 9.3, to obtain:

$$-\text{trace} (X_\alpha - Y_\alpha) \leq f(x_\alpha) - f(y_\alpha) - 2N\delta \rightarrow -2N\delta \quad \text{as } \alpha \rightarrow \infty,$$

contradicting (9.8). This ends the proof of the Lemma. □

Acknowledgements The authors are grateful to D. Bucur for bringing to their attention regularity questions related to the Robin problem, and to J. Manfredi for discussing viscosity solutions. M.L. acknowledges partial support from the NSF grant DMS-1613153. A large part of this work has been completed during M.L. visits to Microsoft Research in Redmond.

Funding M.L. acknowledges partial support from the NSF grant DMS-1613153 and support through visits to MSR (Microsoft Research) in Redmond.

Availability of data and material Not applicable

Code Availability Not applicable

Declarations

Conflict of Interests None.

References

1. Antunovic, T., Peres, Y., Sheffield, S., Somersille, S.: Tug-of-war and infinity laplace equation with vanishing Neumann boundary condition. *Commun. Partial. Differ. Equ.* **37**(10), 1839–1869 (2012)
2. Bass, R., Burdzy, K., Chen, Z.-Q.: On the Robin problem in fractal domains. *Proc. Lond. Math. Soc.* **96**(2), 273–311 (2008)
3. Bass, R., Hsu, P.: Some potential theory for reflecting Brownian motion in Hölder and Lipschitz domains. *Ann. Probab.* **19**(2), 486–508 (1991)
4. Benchérif-Madani, A., Pardoux, É.: A probabilistic formula for a poisson equation with Neumann boundary condition. *Stoch. Anal. Appl.* **27**(4), 739–746 (2009)
5. Brezis, H., Cabrè, X.: Some simple nonlinear PDEs without solutions. *Bollettino dell’Unione Matematica Italiana* **8**(1-B,2), 223–262 (1998)
6. Bucur, D., Giacomini, A., Trebeschi, P.: The Robin-Laplacian problem on varying domains. *Calc. Var.* **55**(6), 55–133 (2016). *Partial Differential Equations*
7. Brosamler, G.A.: A probabilistic solution of the Neumann problem. *Mathematica Scandinavica* **38**(1), 137–147 (1976)
8. Charro, F., Garcia, J., Rossi, J.: A mixed problem for the infinity Laplacian via tug-of-war games. *Calc. Var.* **34**(3), 307–320 (2009)
9. Crandall, M., Ishii, H., Lions, P.-L.: User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.* **27**, 1–67 (1992)
10. Doob, J.L.: *Classical Potential Theory and its Probabilistic Counterpart*. Springer, New York (1984)
11. Gilbarg, D., Trudinger, N.: *Elliptic Partial Differential Equations of Second Order*. *Classics in Mathematics*, 2nd edn. Springer, Berlin Heidelberg (2001)
12. Ishii, H.: On the equivalence of two notions of weak solutions, viscosity solutions and distributional solutions. *Funkcialaj Ekvacioj* **38**, 101–120 (1995)
13. Knothe, H.: Contributions to the theory of convex bodies. *Mich. Math. J.* **4**, 39–52 (1957)

14. Lewicka, M., Manfredi, J., Ricciotti, D.: Random walks and random tug of war in the Heisenberg group. *Mathematische Annalen* **377**, 797–846 (2020)
15. Lewicka, M., Peres, Y.: Which domains have two-sided supporting unit spheres at every boundary point? *Expositiones Mathematicae* **38**(4), 548–558 (2020)
16. Lewicka, M., Peres, Y.: The Robin mean value equation II: asymptotic Hölder regularity submitted (2019)
17. Lieberman, G.: *Oblique Derivative Problems for Elliptic Equations*. World Scientific Publishing (2013)
18. Lucas, K.: Submanifolds of dimension $n - 1$ in E^n with normals satisfying a lipschitz condition, studies in eigenvalue problems. Technical Report 18, Dept of Mathematics, University of Kansas (1957)
19. Manfredi, J., Parviainen, M., Rossi, J.: On the definition and properties of p -harmonious functions. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **11**(2), 215–241 (2012)
20. Morel, J.-M., Oswald, L.: A uniform formulation for hopf maximum principle. Unpublished preprint (1985)
21. Nittka, R.: Quasilinear elliptic and parabolic robin problems on Lipschitz domains. *Nonlinear Differ. Equ. Appl.* **20**(3), 1125–1155 (2013)
22. Papanicolaou, V.: The probabilistic solution of the third boundary value problem for second order elliptic equations. *Probab. Theory Relat. Fields* **87**(1), 27–77 (1990)
23. Peres, Y., Schramm, O., Sheffield, S., Wilson, D.: Tug-of-war and the infinity Laplacian. *J. Amer. Math. Soc.* **22**, 167–210 (2009)
24. Peres, Y., Sheffield, S.: Tug-of-war with noise: A game theoretic view of the p -Laplacian. *Duke Math. J.* **145**(1), 91–120 (2008)
25. Rosenblatt, M.: Remarks on a multivariate transformation. *Ann. Math. Stat.* **23**, 470–472 (1952)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.